

Research Article

Helicity Amplitudes for Generic Multibody Particle Decays Featuring Multiple Decay Chains

Daniele Marangotto 

Università degli Studi di Milano and INFN Milano, Via Celoria 16, 20133 Milano, Italy

Correspondence should be addressed to Daniele Marangotto; daniele.marangotto@unimi.it

Received 4 October 2020; Accepted 30 November 2020; Published 21 December 2020

Academic Editor: Sunny Vagnozzi

Copyright © 2020 Daniele Marangotto. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The publication of this article was funded by SCOAP³.

We present the general expression of helicity amplitudes for generic multibody particle decays characterised by multiple decay chains. This is achieved by addressing for the first time the issue of the matching of the final particle spin states among different decay chains in full generality for generic multibody decays, proposing a method able to match the exact definition of spin states relative to the decaying particle ones. We stress the importance of our result by showing that one of the matching methods used in the literature is incorrect, leading to amplitude models violating rotational invariance. The results presented are therefore relevant for performing numerous amplitude analyses, notably those searching for exotic structures like pentaquarks.

1. Introduction

The helicity formalism, proposed by Jacob and Wick [1] in 1959 to treat relativistic processes involving particles with spin, is still one of the most important tools for performing amplitude analyses of particle decays. To date, complex amplitude analyses involving final-state particles with spin and multiple decay chains have been performed, especially for the search of new resonant structures. Pentaquark searches are a typical example: a pentaquark involves at least one baryon in the final state and introduces an additional decay chain. For instance, pentaquark states were discovered by the LHCb collaboration performing an amplitude analysis of the $\Lambda_b^0 \rightarrow J/\psi p K^-$ baryon decay [2].

However, a consistent definition of the final particle spin states for these kinds of decays turned out to be an issue, since the definition of helicity states is different for different decay chains. Various solutions to match the final particle spin states have been proposed [2–5], but none addressed the problem in full generality for generic multibody decays. In this paper, we present a general method for matching spin states, obtained requiring that, for any decay chain, the final particle states are defined by the same Lorentz transformations relatively to the decaying particle spin states.

To this end, we first review the definition of spin states in quantum mechanics in Section 2, with a particular attention to their phase specifications. The key point we want to stress is that the relative phases among sets of spin states linked by rotations are fully specified by the transformations applied. Therefore, phase differences like those arising when spin states are rotated with respect to their quantisation axis or like the change of sign under 2π angle rotations of fermion states can not be neglected in helicity amplitudes.

Then, we revisit the helicity formalism as originally proposed by Jacob and Wick [1], highlighting the different treatments of daughter particle helicity states in two-body processes. We also propose a simpler definition of two-particle helicity states than the standard one, which allows for an easier matching of the final particle spin states.

In Section 4, we present how to write helicity amplitudes with a consistent definition of the final particle spin states for different decay chains, applicable to any multibody decay topology. We explicitly derive helicity amplitudes for three-body decays.

We stress the need for a consistent definition of the final particle spin states in Section 5. First, we discuss the consequences of an incorrect phase introduced between amplitudes describing different decay chains on the decay distributions,

showing that they produce observable effects on the decay distributions via interference terms. Next, we perform a numerical study on $\Lambda_c^+ \rightarrow pK^-\pi^+$ helicity amplitude models featuring different methods to match the final particle spin states, checking a general property of the decay distributions following from rotational invariance. We show how the method employed for the amplitude analyses [2, 3] is incorrect, leading to amplitude models violating rotational invariance, while that proposed in this article fully satisfies rotational symmetry.

2. On Spin State Definition

In this section, we review the definition of spin states in quantum mechanics underlining the importance of their phase specification, which will be needed for the upcoming discussion of multibody particle decays in the helicity formalism.

In quantum mechanics, the spin of a particle is described by a vector of spin operators $\widehat{\mathbf{S}} = (\widehat{S}_x, \widehat{S}_y, \widehat{S}_z)$, which defines a right-handed spin coordinate system (x, y, z) . The spin states $|s, m\rangle$ are defined as the simultaneous eigenstates of the spin squared modulus $S\Lambda^2$ and \widehat{S}_z , with eigenvalues $s(s+1)$ and m , respectively. The z -axis is called the quantisation axis, while x - and y -axes will be named the orthogonal axes.

The choice of the orthogonal axes specifies the relative phases among spin states, which can be conventionally chosen by defining the action of the ‘‘ladder’’ operators $\widehat{S}_\pm = \widehat{S}_x \pm i\widehat{S}_y$ transforming $|s, m\rangle$ into $|s, m \pm 1\rangle$ eigenstates (see, e.g., Ref. [6]). The overall phase of the spin states is undefined and can be chosen arbitrarily.

Now, let us consider a set of spin states $|s, m\rangle'$ defined relative to the original one $|s, m\rangle$ by applying a rotation \widehat{R} :

$$|s, m\rangle' = \widehat{R}|s, m\rangle. \quad (1)$$

The rotation R defines the relative phases among the two sets: for instance, a rotation around the z -axis of angle α introduces a phase difference between original and rotated spin states:

$$|s, m\rangle' = \widehat{R}_z(\alpha)|s, m\rangle = e^{-i\alpha S\Lambda_z}|s, m\rangle = e^{-i\alpha m}|s, m\rangle. \quad (2)$$

Therefore, once an overall phase for the original set of spin states is conventionally chosen, that of the rotated spin states is defined by the rotation. In other words, the rotated states are completely defined in terms of the original states and the rotation.

The fact that the expectation values of the spin operators transform as a vector under rotations (see, e.g., Ref. [6]) can give the deceptive impression that one can represent rotations applied to spin states in the usual Cartesian space, while spin states transform under spin s representations of the $SU(2)$ group. This is why graphical descriptions are not used in the present article, though they are widely used in the lit-

erature. For instance, it is well known that fermion states change sign for a 2π angle rotation around any axis i :

$$\widehat{R}_i(2\pi)|s, m\rangle = e^{-2i\pi S\Lambda_i}|s, m\rangle = (-1)^{2s}|s, m\rangle, \quad (3)$$

even if the spin operator expectation values do not change.

When considering sets of spin states relatively defined by rotations, it is important to take into account their relative phase differences, since interference effects can make them observable quantities. In this article, we will show how to properly consider the spin state definition in the case of multibody particle decays with different intermediate states in the helicity formalism (Section 4) and the consequences that an incorrect treatment of spin state definitions has on particle decay distributions (Section 5).

3. Helicity Formalism Revisited

In this section, we revisit the helicity formalism [1], developed to overcome problems related to the treatment of spin in relativistic processes. In particular, we highlight the different roles played by the daughter particles in two-body processes and the importance of consistently specifying the definition of their helicity states, including phases. This is an aspect almost neglected so far, which becomes essential for a correct treatment of decays characterised by multiple interfering decay chains. For a clearer treatment of such aspects and a simpler matching of the final particle spin definitions among different decay chains (Section 4), we also propose a different way to express two-particle helicity states, which ease the control of their definitions. A review of the description of relativistic processes involving particles with spin is reported in Appendix A; there, the definition of canonical and helicity states used throughout the article is presented.

The key point underlying the helicity formalism is the invariance of helicity under rotations, exploited to construct two-particle states which are eigenstates of total angular momentum. Indeed, under rotations, both spin states and the momentum expressing their quantisation axis rotate, so that the projection of the particle spin on the momentum is unchanged.

Let us consider a two-body decay $A \rightarrow 1, 2$. The particle 1 helicity states $|\mathbf{p}_1^A, s_1, \lambda_1\rangle$ are defined in the helicity system (see Equation (A.4)):

$$S_1^H = L(-p_1^A \mathbf{z})R(0, -\theta_1, -\phi_1)S_A, \quad (4)$$

with \mathbf{p}_1^A as the particle 1 momentum in the A spin reference rest frame S_A and θ_1, ϕ_1 as its spherical coordinates.

The particle 2 helicity states $|\mathbf{p}_2^A, s_2, \lambda_2\rangle$ are defined in the helicity system:

$$S_2^H = L(-p_2^A \mathbf{z})R(0, -\theta_2, -\phi_2)S_A, \quad (5)$$

which is reached by a different rotation with respect to particle 1 helicity states, so that the direct product of particle 1 and 2 states cannot be related to the total angular momentum S_A .

Exploiting $\mathbf{p}_1^A = -\mathbf{p}_2^A$, in Equations (13) and (14) of Ref. [1], the direct product of daughter particle helicity states is defined as

$$|p_1^A, \theta_1, \phi_1, \lambda_1, \lambda_2\rangle \equiv |\mathbf{p}_1^A, s_1, \lambda_1\rangle \otimes (-1)^{s_2 - \lambda_2} \exp(-i\pi\hat{J}_y) |\mathbf{p}_2^A, s_2, \lambda_2\rangle. \quad (6)$$

These states can be now related to two-particle states with a definite value of total angular momentum, which are denoted $|p_1^A, J, M, \lambda_1, \lambda_2\rangle$ as (here with the $\psi = 0$ convention described in Appendix A)

$$|p_1^A, \theta_1, \phi_1, \lambda_1, \lambda_2\rangle = \sum_{J,M} \sqrt{\frac{2J+1}{4\pi}} D_{M, \lambda_1 - \lambda_2}^J(\phi_1, \theta_1, 0) |p_1^A, J, M, \lambda_1, \lambda_2\rangle. \quad (7)$$

This expression allows writing the $A \rightarrow 1, 2$ decay amplitude as

$$\begin{aligned} \mathcal{A}_{m_A, \lambda_1, \lambda_2}(\theta_1, \phi_1) &= \langle p_1^A, \theta_1, \phi_1, \lambda_1, \lambda_2 | \hat{T} | s_A, m_A \rangle \\ &= \mathcal{H}_{\lambda_1, \lambda_2} D_{m_A, \lambda_1 - \lambda_2}^{*s_A}(\phi_1, \theta_1, 0), \end{aligned} \quad (8)$$

in which $|s_A, m_A\rangle$ are the A spin states defined in the S_A system, \hat{T} is the transition operator, and

$$\mathcal{H}_{\lambda_1, \lambda_2} \equiv \langle J = s_A, M = m_A, \lambda_1, \lambda_2 | \hat{T} | s_A, m_A \rangle \quad (9)$$

are complex numbers called helicity couplings, describing the decay dynamics. Note that the key point underlying Equation (7) is that the state $|p_1^A, 0, 0, \lambda_1, \lambda_2\rangle$ (with \mathbf{p}_1^A aligned with the z -axis) is an eigenstate of \hat{J}_z with eigenvalue $\lambda_1 - \lambda_2$, which is then rotated to

$$|p_1^A, \theta_1, \phi_1, \lambda_1, \lambda_2\rangle = \hat{R}(\phi_1, \theta_1, 0) |p_1^A, 0, 0, \lambda_1, \lambda_2\rangle. \quad (10)$$

We note en passant that, using the properties of Wigner D -matrices (Equation (B.9)), the two-body amplitude can be rewritten as

$$\mathcal{A}_{m_A, \lambda_1, \lambda_2}(\theta_1, \phi_1) = \mathcal{H}_{\lambda_1, \lambda_2} D_{\lambda_1 - \lambda_2, m_A}^{s_A}(0, -\theta_1, -\phi_1), \quad (11)$$

so that, compared with Equation (B.5), the Wigner D -matrix is indeed the representation of the helicity rotation $R(0, -\theta_1, -\phi_1)$ aligning the \mathbf{p}_1^A momentum to the z -axis on the A particle spin states $|s_A, m_A\rangle$.

Let us now consider the particle 2 state in Equation (6): the rotation $\exp(-i\pi\hat{S}_y)$ acts on the helicity state inverting the z -axis direction; therefore, the particle 2 states entering the amplitude (Equation (7)) are actually opposite-helicity states, which represent spin projection eigenstates in the direction opposite to \mathbf{p}_1^A . This different role of particle 1 and 2 states must be properly considered when these particles have a subsequent decay: the amplitude for the particle 2 decay must take into account that it is not referred to $|p_2^A, s_2, \lambda_2\rangle$ states but to those obtained applying the inversion

$\exp(-i\pi\hat{S}_y)$ and the phase factor $(-1)^{s_2 - \lambda_2}$. We stress that these tricky aspects related to the helicity formalism have been neglected or underestimated so far, because for simple processes (like decays via single decay chains or involving spinless particles), they do not have consequences on the decay distributions. However, they matter for the treatment of the more general decays considered in this article.

To take into account in a cleaner way the different roles of particles 1 and 2 in the helicity formalism, we propose a simpler definition of the two-particle state (Equation (6)), which allows for an easier matching of the final particle spin definitions among different decay chains (Section 4). We define the two-particle product state as

$$|p_1^A, \theta_1, \phi_1, \lambda_1, \bar{\lambda}_2\rangle = |\mathbf{p}_1^A, s_1, \lambda_1\rangle \otimes |\mathbf{p}_2^A, s_2, \bar{\lambda}_2\rangle, \quad (12)$$

in which $|\mathbf{p}_2^A, s_2, \bar{\lambda}_2\rangle$ represent spin projection eigenstates in the direction opposite to \mathbf{p}_2^A . The operator $\hat{\bar{\lambda}}$ is the opposite of the helicity:

$$\hat{\bar{\lambda}} = -\hat{\mathbf{S}} \cdot \frac{\mathbf{p}}{p}, \quad (13)$$

and the opposite-helicity reference system of particle 2 is defined by

$$S_2^{OH} = L(p_2^A \mathbf{z}) R(0, -\theta_1, -\phi_1) S_A, \quad (14)$$

that is, the particle 2 rest frame is reached by boosting along its momentum \mathbf{p}_2^A pointing in the direction opposite to the z -axis. Comparing Equations (4) and (14), we see that both particle 1 and particle 2 states are obtained from the same rotation $R(0, -\theta_1, -\phi_1)$, so that their spin is referred to the “same” spin reference system (they only differ by a boost along the z -axis), including the same definition of the orthogonal axes.

It is therefore possible to define eigenstates of total angular momentum $|p_1^A, J, M, \lambda_1, \bar{\lambda}_2\rangle$ similarly as before, and Equation (7) holds with the substitution $-\lambda_2 \rightarrow \bar{\lambda}_2$:

$$|p_1^A, \theta_1, \phi_1, \lambda_1, \bar{\lambda}_2\rangle = \sum_{J,M} \sqrt{\frac{2J+1}{4\pi}} D_{M, \lambda_1 + \bar{\lambda}_2}^J(\phi_1, \theta_1, 0) |p_1^A, J, M, \lambda_1, \bar{\lambda}_2\rangle. \quad (15)$$

The two-body decay amplitude becomes

$$\begin{aligned} \mathcal{A}_{m_A, \lambda_1, \bar{\lambda}_2}(\theta_1, \phi_1) &= \langle p_1^A, \theta_1, \phi_1, \lambda_1, \bar{\lambda}_2 | \hat{T} | s_A, m_A \rangle \\ &= \mathcal{H}_{\lambda_1, \bar{\lambda}_2} D_{m_A, \lambda_1 + \bar{\lambda}_2}^{*s_A}(\phi_1, \theta_1, 0), \end{aligned} \quad (16)$$

and the helicity values allowed by angular momentum conservation are

$$|\lambda_1| \leq s_1, |\bar{\lambda}_2| \leq s_2, |\lambda_1 + \bar{\lambda}_2| \leq s_A. \quad (17)$$

The amplitudes (Equations (8) and (16)) are the same but for the substitution $\lambda_2 \leftrightarrow \bar{\lambda}_2$, so why bother with a new state definition? The difference is in the definition of

particle 2 states: in the standard formulation (Equation (6)), the particle 2 opposite-helicity state is obtained inverting the helicity one, applying two rotations to the initial system S_A plus a phase; in our definition, it is just defined by a single rotation from S_A . Our choice simplifies both the writing of particle 2 subsequent decay amplitudes and the matching of the final particle spin states among different decay chains.

For the purpose of Section 4, it is useful to derive the relation between opposite-helicity and canonical states, the analogue of Equation (A.6) for helicity states. It is obtained applying Equation (A.5) along with the relations $\mathbf{p}_2^A = -\mathbf{p}_1^A$, $p_2^A = p_1^A$, to the definition of canonical states (Equation (A.1)):

$$\begin{aligned} S_2^C &= L(-\mathbf{p}_2^A)S_A = L(\mathbf{p}_1^A)S_A = R(\phi_1, \theta_1, 0)L(p_1^A \mathbf{Z})R(0, -\theta_1, -\phi_1)S_A \\ &= R(\phi_1, \theta_1, 0)L(p_2^A \mathbf{Z})R(0, -\theta_1, -\phi_1)S_A = R(\phi_1, \theta_1, 0)S_2^{OH}. \end{aligned} \quad (18)$$

The rotation is indeed the same as the one from the helicity to the canonical system of particle 1 (see Equation (A.6)):

$$S_1^C = R(\phi_1, \theta_1, 0)S_1^H. \quad (19)$$

4. Helicity Amplitudes for Generic Multibody Particle Decays Featuring Multiple Decay Chains

In this section, we present how helicity amplitudes for generic multibody particle decays characterised by multiple decay chains can be written: in particular, we propose an original method to match the final particle spin states among different decay chains able to properly take into account the definition of spin states. For the sake of clarity, we consider a three-body decay $A \rightarrow 1, 2, 3$, but the method presented to write helicity amplitudes is applicable to any decay topology.

Decay amplitudes for multibody particle decays are obtained in the helicity formalism by breaking the decay chain in sequential two-body decays mediated by intermediate states; for instance, a three-body decay is treated by breaking it into two binary decays. Three decay chains, involving three kinds of intermediate states, are possible: $A \rightarrow R(\rightarrow 1, 2), 3$, $A \rightarrow S(\rightarrow 1, 3), 2$, and $A \rightarrow U(\rightarrow 2, 3), 1$.

We first consider the $A \rightarrow R(\rightarrow 1, 2), 3$ decay chain: the $A \rightarrow R, 3$ decay can be expressed in the A rest frame by Equation (16):

$$\begin{aligned} \mathcal{A}_{m_A, \lambda_R, \bar{\lambda}_3}^{A \rightarrow R, 3}(\theta_R, \phi_R) &= \langle p_1^A, \theta_R, \phi_R, \lambda_R, \bar{\lambda}_3^R | \hat{T} | s_A, m_A \rangle \\ &= \mathcal{H}_{\lambda_R, \bar{\lambda}_3}^{A \rightarrow R, 3} D_{m_A, \lambda_R + \bar{\lambda}_3}^{*s_A}(\phi_R, \theta_R, 0), \end{aligned} \quad (20)$$

and the $R \rightarrow 1, 2$ decay can be written in the same form, in the R rest frame, by applying Equation (16) to the R state $|s_R, \lambda_R\rangle$ as the decaying particle:

$$\begin{aligned} \mathcal{A}_{\lambda_R, \lambda_1^R, \bar{\lambda}_2^R}^{R \rightarrow 1, 2}(\theta_1^R, \phi_1^R) &= \langle p_1^R, \theta_1^R, \phi_1^R, \lambda_1^R, \bar{\lambda}_2^R | \hat{T} | \mathbf{p}_R^A, s_R, \lambda_R \rangle \\ &= \mathcal{H}_{\lambda_1^R, \bar{\lambda}_2^R}^{R \rightarrow 1, 2} D_{m_R, \lambda_1^R + \bar{\lambda}_2^R}^{*s_R}(\phi_1^R, \theta_1^R, 0). \end{aligned} \quad (21)$$

The R superscript is put on helicity values and angles of particles 1 and 2 to stress that their definition is specific to the $A \rightarrow R(\rightarrow 1, 2), 3$ decay chain.

The total amplitude of the $A \rightarrow R(\rightarrow 1, 2), 3$ decay is written introducing R as the intermediate state, and summing the amplitudes over the helicity values λ_R satisfies the angular momentum conservation requirements (Equation (17)):

$$\begin{aligned} \mathcal{A}_{m_A, \lambda_1^R, \bar{\lambda}_2^R, \bar{\lambda}_3^R}^{A \rightarrow R, 3 \rightarrow 1, 2, 3}(\Omega) &= \langle \{\mathbf{p}_i\}, \lambda_1^R, \bar{\lambda}_2^R, \bar{\lambda}_3^R | \hat{T} | s_A, m_A \rangle \\ &= \sum_{\lambda_R} \langle p_1^R, \theta_1^R, \phi_1^R, \lambda_1^R, \bar{\lambda}_2^R | \hat{T} | \mathbf{p}_R^A, s_R, \lambda_R \rangle \\ &\quad \times \langle p_1^A, \theta_R, \phi_R, \lambda_R, \bar{\lambda}_3^R | \hat{T} | s_A, m_A \rangle \\ &= \sum_{\lambda_R} \mathcal{A}_{m_A, \lambda_R, \bar{\lambda}_3}^{A \rightarrow R, 3}(\theta_R, \phi_R) \mathcal{A}_{\lambda_R, \lambda_1^R, \bar{\lambda}_2^R}^{R \rightarrow 1, 2}(\theta_1^R, \phi_1^R) \\ &= \sum_{\lambda_R} \mathcal{H}_{\lambda_1^R, \bar{\lambda}_2^R}^{R \rightarrow 1, 2} D_{\lambda_R, \lambda_1^R + \bar{\lambda}_2^R}^{*s_R}(\phi_1^R, \theta_1^R, 0) \\ &\quad \times \mathcal{H}_{\lambda_R, \bar{\lambda}_3}^{A \rightarrow R, 3} D_{m_A, \lambda_R + \bar{\lambda}_3}^{*s_A}(\phi_R, \theta_R, 0). \end{aligned} \quad (22)$$

Note that the angles entering the decay amplitude depend on the phase space variables describing the decay, denoted collectively as Ω .

Now, let us consider the $A \rightarrow S(\rightarrow 1, 3), 2$ decay chain. Its associated amplitude is, following Equation (22),

$$\begin{aligned} \mathcal{A}_{m_A, \lambda_1^S, \bar{\lambda}_2^S, \bar{\lambda}_3^S}^{A \rightarrow S, 2 \rightarrow 1, 2, 3}(\Omega) &= \sum_{\lambda_S} \mathcal{H}_{\lambda_1^S, \bar{\lambda}_2^S}^{S \rightarrow 1, 3} D_{\lambda_S, \lambda_1^S + \bar{\lambda}_2^S}^{*s_S}(\phi_1^S, \theta_1^S, 0) \\ &\quad \times \mathcal{H}_{\lambda_S, \bar{\lambda}_3^S}^{A \rightarrow S, 2} D_{m_A, \lambda_S + \bar{\lambda}_3^S}^{*s_A}(\phi_S, \theta_S, 0). \end{aligned} \quad (23)$$

Helicity values and angles denoted by the S superscripts are defined specifically for the $A \rightarrow S(\rightarrow 1, 3), 2$ decay chain: the definition of the final particle spin states for this decay chain is different from that used for the R intermediate state one.

To write the total amplitude of the $A \rightarrow 1, 2, 3$ decay, amplitudes associated with different intermediate states must be summed coherently to properly include interference effects. The summing can be performed only if the definition of the final particle spin states is the same across different decay chains. Since helicity systems are specific to each decay chain, they must be rotated to a reference set of spin states, for each final particle. Various solutions to match the final particle spin states have been proposed [2–5], but none addressed the problem in full generality for generic multibody decays. In the following, we derive the correct matching of the final particle spin states requiring that, for any decay chain, the final particle states are defined by the same Lorentz transformations relatively to the decaying particle spin states.

The definition of the helicity states used to express the helicity amplitudes (Equations (22) and (23)) is given relatively to the A particle spin states (S_A reference system) by a sequence of Lorentz transformations. Once a conventional definition of the $|s_A, m_A\rangle$ state overall phase is chosen (see

Section 2), the helicity states are fully specified by the Lorentz transformation sequence, overall phase included. Therefore, to relate different helicity state definitions, it is mandatory to refer back to the initial S_A reference system; i.e., it is not possible to relate the two systems via a direct transformation. To stress this essential point, let us consider the two helicity systems for particle 1 defined by the R and S decay chains, $S_1^{HH,R}$ and $S_1^{HH,S}$, respectively. Suppose we find a rotation \widehat{R} such that $\widehat{R}S_1^{HH,S} = S_1^{HH,R}$ and we rotate the $|\mathbf{p}_1^S, s_1, \lambda_1^S\rangle$ states applying the Wigner D -matrix associated with that rotation. However, this does not guarantee that the spin state phase definition is the same between $S_1^{HH,R}$ and $\widehat{R}S_1^{HH,S}$ systems, since they are defined with respect to S_A by different Lorentz transformation sequences: for a fermion, the two may differ by an overall 2π rotation changing the relative sign among spin states by Equation (3).

The correct way to proceed is to define a reference spin system for the final particle from the initial system S_A and relate each helicity system to this one by applying a sequence of rotations which turn the Lorentz transformation sequence defining the helicity state into that specifying the reference one. The reference and the helicity systems are different rest frames of the final particle, so no boosts need to be applied. Any spin system can be chosen as the reference one, and we will choose canonical states reached from S_A being the simplest possibility.

We illustrate the method explicitly deriving the rotation sequences transforming the helicity systems of the $A \rightarrow 1, 2, 3$ decay final particles, for R and S intermediate state decay chains, to their canonical systems reached from S_A . Starting from particle 1, the helicity system $S_1^{HH,R}$ is defined from S_A by the sequence (see Equation (A.4)):

$$S_R^H = L(-p_R^A \mathbf{z}) R(0, -\theta_R, -\phi_R) S_A, \quad (24)$$

$$S_1^{HH,R} = L(-p_1^R \mathbf{z}) R(0, -\theta_1^R, -\phi_1^R) S_R^H, \quad (25)$$

in which S_R^H is the R helicity system reached from S_A . The canonical system of particle 1 derived from S_R^H , following Equation (A.1) by decomposing the Lorentz boost, is

$$S_1^{HC,R} = R(\phi_1^R, \theta_1^R, 0) L(-p_1^R \mathbf{z}) R(0, -\theta_1^R, -\phi_1^R) S_R^H = R(\phi_1^R, \theta_1^R, 0) S_1^{HH,R}, \quad (26)$$

so that the rotation $R(\phi_1^R, \theta_1^R, 0)$ from $S_1^{HH,R}$ to $S_1^{HC,R}$ systems “undoes” the helicity rotation $R(0, -\theta_1^R, -\phi_1^R)$, acting in the particle 1 rest frame. The spin states $|\mathbf{p}_1^R, s_1, \mu_1^R\rangle$ defined by the $S_1^{HC,R}$ system Lorentz transformation are expressed in terms of the helicity states $|\mathbf{p}_1^R, s_1, \lambda_1^R\rangle$ as in Equation (A.7):

$$\widehat{R}(\phi_1^R, \theta_1^R, 0) |\mathbf{p}_1^R, s_1, \lambda_1^R\rangle = \sum_{\mu_1^R} D_{\mu_1^R, \lambda_1^R}^{s_1}(\phi_1^R, \theta_1^R, 0) |\mathbf{p}_1^R, s_1, \mu_1^R\rangle. \quad (27)$$

Now, let us consider the canonical system of particle 1 $S_1^{CC,R}$ derived from the R canonical system S_R^C , defined following Equation (A.1):

$$S_R^C = L(-p_R^A) S_A = R(\phi_R, \theta_R, 0) L(-p_R^A \mathbf{z}) R(0, -\theta_R, -\phi_R) S_A, \quad (28)$$

$$S_1^{CC,R} = L(-p_1^R) S_R^C = R(\phi_1^R, \theta_1^R, 0) L(-p_1^R \mathbf{z}) R(0, -\theta_1^R, -\phi_1^R) S_R^C, \quad (29)$$

in which primed quantities indicate that the particle 1 momentum used to build the $S_1^{CC,R}$ system is different from the one defining the $S_1^{HC,R}$ system, being determined from S_R^C instead of S_R^H . The momenta \mathbf{p}_1^R and \mathbf{p}_1^R differ by the additional rotation $R(\phi_R, \theta_R, 0)$ used in the definition of S_R^C :

$$\mathbf{p}_1^R = R(\phi_R, \theta_R, 0) \mathbf{p}_1^R, \quad (30)$$

and the rotation $R(0, -\theta_1^R, -\phi_1^R)$ used to align \mathbf{p}_1^R to the z -axis can be decomposed into $R(0, -\theta_R, -\phi_R)$, rotating \mathbf{p}_1^R into \mathbf{p}_1^R , times $R(0, -\theta_1^R, -\phi_1^R)$, with the helicity rotation aligning \mathbf{p}_1^R with the z -axis:

$$R(0, -\theta_1^R, -\phi_1^R) = R(0, -\theta_1^R, -\phi_1^R) R(0, -\theta_R, -\phi_R). \quad (31)$$

Applying the above decomposition, along with its inverse,

$$R(\phi_1^R, \theta_1^R, 0) = R(\phi_R, \theta_R, 0) R(\phi_1^R, \theta_1^R, 0), \quad (32)$$

to Equation (29) and using the systems defined in Equations (25) and (26), we obtain

$$\begin{aligned} S_1^{CC,R} &= L(-\mathbf{p}_1^R) L(-\mathbf{p}_R^A) S_A \\ &= R(\phi_1^R, \theta_1^R, 0) L(-p_1^R \mathbf{z}) R(0, -\theta_1^R, -\phi_1^R) \\ &\quad \times R(\phi_R, \theta_R, 0) L(-p_R^A \mathbf{z}) R(0, -\theta_R, -\phi_R) S_A \\ &= R(\phi_1^R, \theta_1^R, 0) L(-p_1^R \mathbf{z}) R(0, -\theta_1^R, -\phi_1^R) \\ &\quad \times L(-p_R^A \mathbf{z}) R(0, -\theta_R, -\phi_R) S_A \\ &= R(\phi_R, \theta_R, 0) R(\phi_1^R, \theta_1^R, 0) S_1^{HH,R} = R(\phi_R, \theta_R, 0) S_1^{HC,R}. \end{aligned} \quad (33)$$

The rotation $R(\phi_R, \theta_R, 0)$ from $S_1^{HC,R}$ to $S_1^{CC,R}$ systems “undoes” the helicity rotation $R(0, -\theta_R, -\phi_R)$, still acting in the particle 1 rest frame. The spin states $|\mathbf{p}_1^R, s_1, \nu_1^R\rangle$ defined




FIGURE 1: Graphical summary of the spin state definitions associated with the decay amplitude for final particle 1 and intermediate state R . The solid lines indicate the Lorentz transformation sequence involved in the writing of the helicity amplitude. Once a conventional choice for $|s_A, m_A\rangle$ initial A particle spin states is set, all the other spin states are completely specified by their Lorentz transformation sequence. Momentum labels of spin states have been suppressed to avoid cluttering the notation.

by the $S_1^{CC,R}$ system are expressed in terms of the states $|\mathbf{p}_1^R, s_1, \mu_1^R\rangle$ as

$$\widehat{R}(\phi_R, \theta_R, 0) |\mathbf{p}_1^R, s_1, \mu_1^R\rangle = \sum_{\nu_1^R} D_{\nu_1^R, \mu_1^R}^{s_1}(\phi_R, \theta_R, 0) |\mathbf{p}_1^R, s_1, \nu_1^R\rangle. \quad (34)$$

Finally, the system $S_1^{CC,R}$ has to be related to our reference system, the canonical system of particle 1 reached directly from S_A :

$$S_1^{C,A} = L(-\mathbf{p}_1^A) S_A, \quad (35)$$

which differs from $S_1^{CC,R}$ by

$$S_1^{C,A} = L(-\mathbf{p}_1^A) L(\mathbf{p}_R^A) L(\mathbf{p}_1^R) S_1^{CC,R}. \quad (36)$$

The transformation $L(-\mathbf{p}_1^A) L(\mathbf{p}_R^A) L(\mathbf{p}_1^R)$ is equivalent to a rotation, called the Wigner rotation, indicated generically as a Euler rotation $R(\alpha_1^{W,R}, \beta_1^{W,R}, \gamma_1^{W,R})$, in which the associated Euler angles do not have a simple algebraic expression but can be easily computed. The value of the Euler angles can be calculated as follows: the Wigner rotation matrix is found from multiplying the three matrices associated with each Lorentz boost; the unit vectors describing the rotated system are simply the columns of the Wigner rotation matrix; finally, Euler angles are computed using Equation (B.4). Explicit axis-angle expressions for the composition of two Lorentz boosts are reported in Ref. [7]. The canonical spin states $|\mathbf{p}_1^A, s_1, m_1\rangle$ defined by the $S_1^{C,A}$ coordinate systems are expressed in terms of the states $|\mathbf{p}_1^R, s_1, \nu_1^R\rangle$ as

$$\begin{aligned} \widehat{R}(\alpha_1^{W,R}, \beta_1^{W,R}, \gamma_1^{W,R}) |\mathbf{p}_1^R, s_1, \nu_1^R\rangle \\ = \sum_{m_1} D_{m_1, \nu_1^R}^{s_1}(\alpha_1^{W,R}, \beta_1^{W,R}, \gamma_1^{W,R}) |\mathbf{p}_1^A, s_1, m_1\rangle. \end{aligned} \quad (37)$$

The different Lorentz transformations applied starting from the initial S_A system down to the final $S_1^{C,A}$ one, for R intermediate states, are graphically summarised in Figure 1, along with the definition of the many spin states involved in the writing of the helicity amplitude.

Combining Equations (27), (34), and (37), the relation between particle 1 canonical spin states defined by the $S_1^{C,A}$ system and particle 1 helicity states defined by the $S_1^{HH,R}$ system, those expressing the decay amplitude (Equation (22)), is

$$\begin{aligned} \widehat{R}(\alpha_1^{W,R}, \beta_1^{W,R}, \gamma_1^{W,R}) \widehat{R}(\phi_R, \theta_R, 0) \widehat{R}(\phi_1^R, \theta_1^R, 0) |\mathbf{p}_1^R, s_1, \lambda_1^R\rangle \\ = \sum_{\mu_1^R} D_{\mu_1^R, \lambda_1^R}^{s_1}(\phi_1^R, \theta_1^R, 0) \times \sum_{\nu_1^R} D_{\nu_1^R, \mu_1^R}^{s_1}(\phi_R, \theta_R, 0) \sum_{m_1} D_{m_1, \nu_1^R}^{s_1} \\ \times (\alpha_1^{W,R}, \beta_1^{W,R}, \gamma_1^{W,R}) |\mathbf{p}_1^A, s_1, m_1\rangle. \end{aligned} \quad (38)$$

The transformation sequence for particle 2 is almost identical to that for particle 1, but for the Lorentz boost involving its momentum, thanks to the use of the opposite-helicity states we introduced in Section 3. The sequence of helicity rotations is indeed the same as for particle 1, since particle 2 opposite-helicity states are defined by the same helicity rotation specifying particle 1 helicity states (see Equation (18)). The relation between particle 2 canonical spin states ($S_2^{C,A}$ system) and particle 2 opposite-helicity states ($S_2^{OHH,S}$ system) is thus the same as Equation (38) apart from the Wigner rotation:

$$R(\alpha_2^{W,R}, \beta_2^{W,R}, \gamma_2^{W,R}) = L(-\mathbf{p}_2^A) L(\mathbf{p}_R^A) L(\mathbf{p}_2^R). \quad (39)$$

For particle 2, the advantage of using our proposed definition for two-particle product helicity states (Equation (12)) instead of the usual one (Equation (6)) is evident: in the latter case, two rotations must be performed, one corresponding to the helicity rotation and one for the inversion.

For particle 3, the spin rotation to its canonical system is much simpler since only one helicity rotation has to be “undone,” with no Wigner rotation needed since a single direct boost from particle A to particle 3 rest frames is involved, leading to

$$|\mathbf{p}_3^A, s_3, \lambda_3^R\rangle = \sum_{m_3} D_{m_3, \lambda_3^R}^{s_3}(\phi_R, \theta_R, 0) |\mathbf{p}_3^A, s_3, m_3\rangle. \quad (40)$$

The final particle spin rotations are introduced in the decay amplitudes as follows. The amplitude for R intermediate states we are interested in is

$$\mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow R, 3 \rightarrow 1, 2, 3}(\Omega) = \langle \{\mathbf{p}_i\} m_1, m_2, m_3 | \widehat{T} | s_A, m_A \rangle, \quad (41)$$

in which the final particles state

$$|\{\mathbf{p}_i\}, m_1, m_2, m_3\rangle = |s_1, m_1\rangle \otimes |s_2, m_2\rangle \otimes |s_3, m_3\rangle \quad (42)$$

is the product of the canonical spin states reached from the S_A system for each final particle.

For the sake of clarity, we consider a much simpler example: the case of a single rotation on a single particle state. Suppose we know the transition amplitude $\mathcal{A}_\lambda = \langle s, \lambda | \widehat{T} | i \rangle$ between a given initial state $|i\rangle$ and a final state $|s, \lambda\rangle$, but we would like to express the amplitude with respect to some rotated spin states:

$$\widehat{R}(\alpha, \beta, \gamma) |s, \lambda\rangle = \sum_m D_{m, \lambda}^s(\alpha, \beta, \gamma) |s, m\rangle. \quad (43)$$

To this end, we write the transition amplitude for the rotated states introducing the $|s, \lambda\rangle$ states as a set of intermediate states:

$$\begin{aligned} \mathcal{A}_m &= \langle s, m | \widehat{T} | i \rangle = \sum_\lambda \langle s, m | \widehat{R}(\alpha, \beta, \gamma) |s, \lambda\rangle \langle s, \lambda | \widehat{T} | i \rangle \\ &= \sum_\lambda D_{m, \lambda}^s(\alpha, \beta, \gamma) \mathcal{A}_\lambda, \end{aligned} \quad (44)$$

in which the rotation operator $\widehat{R}(\alpha, \beta, \gamma)$ acts as a transition operator between the final particle spin states. Note the parallel with Equation (11): the Wigner D -matrix is indeed the representation of the rotation applied.

Generalising the case of a single rotation for a single particle, the amplitude for R intermediate states (Equation (41)), expressed for the final particle canonical state (Equation (42)) in terms of the amplitude for helicity states (Equation (22)), becomes

$$\begin{aligned} \mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow R, 3 \rightarrow 1, 2, 3}(\Omega) &= \sum_{\lambda_1^R, \mu_1^R, \nu_1^R} D_{m_1, \nu_1^R}^{s_1}(\alpha_1^{W, R}, \beta_1^{W, R}, \gamma_1^{W, R}) D_{\nu_1^R, \mu_1^R}^{s_1} \\ &\cdot (\phi_R, \theta_R, 0) D_{\mu_1^R, \lambda_1^R}^{s_1}(\phi_1^R, \theta_1^R, 0) \\ &\times \sum_{\lambda_2^R, \mu_2^R, \nu_2^R} D_{m_2, \nu_2^R}^{s_2}(\alpha_2^{W, R}, \beta_2^{W, R}, \gamma_2^{W, R}) D_{\nu_2^R, \mu_2^R}^{s_2} \\ &\cdot (\phi_R, \theta_R, 0) D_{\mu_2^R, \lambda_2^R}^{s_2}(\phi_1^R, \theta_1^R, 0) \\ &\times \sum_{\lambda_3^R} D_{m_3, \lambda_3^R}^{s_3}(\phi_R, \theta_R, 0) \times \mathcal{A}_{m_A, \lambda_1^R, \lambda_2^R, \lambda_3^R}^{A \rightarrow R, 3 \rightarrow 1, 2, 3}(\Omega). \end{aligned} \quad (45)$$

The amplitude for S intermediate states can be written analogously, taking into account the different decay topologies (interchanging particle 2 and 3 roles):

$$\begin{aligned} \mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow S, 2 \rightarrow 1, 2, 3}(\Omega) &= \sum_{\lambda_1^S, \mu_1^S, \nu_1^S} D_{m_1, \nu_1^S}^{s_1}(\alpha_1^{W, S}, \beta_1^{W, S}, \gamma_1^{W, S}) D_{\nu_1^S, \mu_1^S}^{s_1} \\ &\cdot (\phi_S, \theta_S, 0) D_{\mu_1^S, \lambda_1^S}^{s_1}(\phi_1^S, \theta_1^S, 0) \\ &\times \sum_{\lambda_2^S} D_{m_2, \lambda_2^S}^{s_2}(\phi_S, \theta_S, 0) \times \sum_{\lambda_3^S, \mu_3^S, \nu_3^S} D_{m_3, \nu_3^S}^{s_3} \\ &\cdot (\alpha_3^{W, S}, \beta_3^{W, S}, \gamma_3^{W, S}) D_{\nu_3^S, \mu_3^S}^{s_3}(\phi_S, \theta_S, 0) D_{\mu_3^S, \lambda_3^S}^{s_3} \\ &\cdot (\phi_1^S, \theta_1^S, 0) \times \mathcal{A}_{m_A, \lambda_1^S, \lambda_2^S, \lambda_3^S}^{A \rightarrow S, 2 \rightarrow 1, 2, 3}(\Omega). \end{aligned} \quad (46)$$

Finally, the total decay amplitude is obtained summing the amplitudes associated with each intermediate state:

$$\begin{aligned} \mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow 1, 2, 3}(\Omega) &= \sum_i \mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow R_i, 3 \rightarrow 1, 2, 3}(\Omega) + \sum_j \mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow S_j, 2 \rightarrow 1, 2, 3}(\Omega) \\ &+ \sum_k \mathcal{A}_{m_A, m_1, m_2, m_3}^{A \rightarrow U_k, 1 \rightarrow 1, 2, 3}(\Omega), \end{aligned} \quad (47)$$

in which the amplitude for U intermediate states can be written similarly as for R and S states.

For a generic multibody decay, the generalisation of the method for the matching of the final particle spin states is straightforward. Each final particle helicity state must be rotated “undoing” all the helicity rotations applied along its decay chain, in reversed order. One additional Wigner rotation is needed, describing the difference between the direct boost defining the particle canonical state and the boost sequence applied along its decay chain.

5. Effects of an Incorrect Matching of the Final Particle Spin States on Decay Distributions

We stress the need for a correct matching of the final particle spin states by considering the consequences an incorrect matching can have. We first present a general discussion of the observable effects one can introduce in the decay distributions, and even in the simple case, the incorrect matching just introduces a relative phase among spin states belonging to different decay chains. Then, we present a numerical study in which we compare the method for the matching of the final particle spin states presented in this article to the one employed in Refs. [2, 3]: we show that the latter is not correct since it breaks rotational invariance.

We first discuss the general case of a particle A decay to $\{i = 1, \dots, n\}$ final state particles, passing through two intermediate states R and S . The associated amplitudes are denoted as $\mathcal{A}_{m_A, \{m_i\}}^R$ and $\mathcal{A}_{m_A, \{m_i\}}^S$. Let us suppose that, due to an incorrect matching of the final particle spin state definition, a phase difference $\exp[i\psi(\Omega)]$ is introduced between the two amplitudes. Since the definition of the spin systems depends on the phase space variables, the incorrect phase will

be in general function of the decay phase space. The effects of such a phase are in a certain sense analogous to the phase effects seen in interferometry experiments: in that case, the phase difference between particles following different paths is a physical effect produced by a difference in energy potential felt by the particle; in our case, it is an unnatural effect caused by an incorrect definition of spin states among different decay chains (the analogue of paths). Also, the mere change of sign of fermion states under 2π angle rotations is observable, being measured in neutron interferometry [8, 9].

The polarised decay rate for definite initial m_A and final spin projections $\{m_i\}$ (see Equation (B.5) of Appendix C) is

$$\begin{aligned} P_{m_A, \{m_i\}}(\Omega) &= \left| \mathcal{A}_{m_A, \{m_i\}}^R + \exp[i\psi(\Omega)] \mathcal{A}_{m_A, \{m_i\}}^S \right|^2 \\ &= \left| \mathcal{A}_{m_A, \{m_i\}}^R \right|^2 + \left| \mathcal{A}_{m_A, \{m_i\}}^S \right|^2 \\ &\quad + 2 \operatorname{Re} \left[\exp[-i\psi(\Omega)] \mathcal{A}_{m_A, \{m_i\}}^R \mathcal{A}_{m_A, \{m_i\}}^{S*} \right]. \end{aligned} \quad (48)$$

The incorrect phase affects the decay rate modifying the interference terms between different decay chains, changing their functional form in terms of phase space variables. In an amplitude fit to experimental data, it means that fit parameters depending on interference effects between different decay chains can be biased due to the incorrect phase space dependence introduced. Note that the incorrect phase also affects the unpolarised decay rate: there is no guarantee that the sum over the spin states mitigates its effect.

A numerical study illustrating these kinds of unnatural effects is performed on the decay distributions of the three-body $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay obtained from its helicity amplitudes. The three-body decay phase space is described by 5 degrees of freedom: two ‘‘Dalitz’’ two-body invariant masses and three angles describing the orientation of the decay with respect to the A reference system S_A , called orientation angles in the following. For the $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay, they can be chosen to be $m_{pK^-}^2$, $m_{K^-\pi^+}^2$, the cosine of the proton polar angle in the $S_{\Lambda_c^+}$ system, $\cos \theta_p$, the proton azimuthal angle, ϕ_p , and the signed angle between the plane formed by the proton and the Λ_c^+ quantisation axis and the plane formed by the kaon and the pion, named χ .

We test a property of the decay distributions following from rotational invariance, which must be satisfied irrespective of the amplitude model considered: for zero Λ_c^+ polarisation, the orientation angle distributions $\cos \theta_p$, ϕ_p , χ must be uniform. Indeed, in the absence of a polarisation vector, nothing specifies a direction in the $S_{\Lambda_c^+}$ system. This property provides a necessary test for the correctness of an amplitude model: if the model produces anisotropic orientation angle distributions for zero polarisation, it is wrong, violating rotational invariance.

We consider $\Lambda_c^+ \rightarrow pK^-\pi^+$ helicity amplitudes written closely following the three-body ones described in Section 4 (details specific to the $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay are reported in Appendix D) applying two different methods for matching proton spin states among different decay chains: the one pre-

sented in this article and the one employed for the amplitude analyses [2, 3]. For the $\Lambda_c^+ \rightarrow pK^-\pi^+$ case, the latter method consists in applying a single rotation to the proton states aligning their quantisation axes to a reference one, which are defined as the direction opposite to the momentum of the particle recoiling against the proton in the proton rest frame. Taking the $K^* (\rightarrow K^-\pi^+)$ decay chain as the reference, the rotation applied to the proton helicity states is

$$\widehat{R}(0, \beta_p^R, 0) \left| \mathbf{p}_p^R, 1/2, \lambda_p \right\rangle = d_{\lambda_p}^{1/2}(\beta_p^R) \left| \mathbf{p}_p^R, 1/2, \lambda_p' \right\rangle, \quad (49)$$

with angles

$$\cos \beta_p^{\Lambda^*} = \widehat{\mathbf{p}}_{K^*}^p \cdot \widehat{\mathbf{p}}_{K^-}^p, \quad \cos \beta_p^{\Delta^*} = \widehat{\mathbf{p}}_{K^*}^p \cdot \widehat{\mathbf{p}}_{\pi^+}^p, \quad (50)$$

for the $\Lambda^* (\rightarrow pK^-)$ and $\Delta^{*++} (\rightarrow p\pi^+)$ decay chains, respectively.

For the numerical study, we consider a $\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude model consisting of three resonance states, one per decay channel. The detailed specification of the helicity couplings and resonance descriptions employed is reported in Appendix D. The code reproducing the amplitude model is based on a version of the TensorFlowAnalysis package [10] adapted to five-dimensional phase space three-body amplitude fits [11]; this package depends on the machine learning framework TensorFlow [12] and the ROOT package [13].

The phase space distributions are described by a set of five millions Monte Carlo pseudodata generated according to the two amplitude models for zero polarisation. We stress that the two amplitude models only differ by the proton spin matching method.

The distributions associated with the method presented in this article are shown in Figure 2: the orientation angle distributions are precisely isotropic, given the accuracy allowed by the large pseudodata sample.

The distributions associated with the spin matching method used in Refs. [2, 3] are shown in Figure 3: the orientation angle distributions are evidently anisotropic, also featuring a step in the ϕ_p distribution. This method is thus incorrect, leading to an amplitude model violating rotational invariance. Moreover, it is important to note that the invariant mass distributions are different for the two methods: therefore, an incorrect spin matching can also affect Dalitz plot analyses, in which orientation angles are integrated over the decay rate, as in spin-zero meson decays or when the polarisation of the decaying particle is not considered.

To conclude, we discuss why the matching method of Refs. [2, 3] is not correct. The proton spin rotations (Equation (49)) ensure that the definition of the proton quantisation axis is the same among different decay chains. Therefore, the proton states can differ among different decay chains by a rotation of the orthogonal axes (Equation (2)) or by the sign of the proton state (Equation (3)), which introduce an incorrect phase difference among decay chains. This is exactly the case considered at the beginning of this section. Indeed, the decay distributions associated with single resonances are the same for the two matching methods (up to




FIGURE 2: Phase space decay distributions for the $\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude models written using the proton spin matching method presented in this article. “Model” refers to the full $\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude model distributions, while “D,” “L,” and “K” labels indicate the distributions associated with single Δ^{*++} , Λ^* , K^* resonance contributions.

an irrelevant normalisation factor) (Figure 3), since the phase difference $\exp[i\psi(\Omega)]$ cancels for single resonance decay rates. However, the incorrect phase affects the decay distributions of the full amplitude model, which includes interference effects among different decay chains, as in Equation (48).

6. Conclusions

We presented the general expression of helicity amplitudes for generic multibody particle decays characterised by multiple decay chains. We demonstrated the importance of a precise specification of spin states for correct writing of helicity




FIGURE 3: Phase space decay distributions for the $\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude models written using the proton spin matching method used in Refs. [2, 3]. “Model” refers to the full $\Lambda_c^+ \rightarrow pK^-\pi^+$ amplitude model distributions, while “D,” “L,” and “K” labels indicate the distributions associated with single Δ^{*++} , Λ^* , K^* resonance contributions.

amplitudes, proposing an original method to match the final particle spin states among different decay chains, applicable to any multibody decay topology.

To this end, we reviewed the definition of spin states in quantum mechanics, with a particular attention to their phase specifications, and the helicity formalism, considering a simpler definition of two-particle helicity states than

the standard one. The proposed method to match the final particle spin states was obtained requiring that, for any decay chain, the final particle states are defined by the same Lorentz transformations relatively to the decaying particle spin states, by applying a sequence of rotations. Helicity amplitudes were explicitly written for three-body decays.

We discussed the consequences of an incorrect phase introduced between amplitudes describing different intermediate states on the decay distributions, showing that they produce observable effects on the decay distributions via interference terms. We also tested numerically our spin matching method against the one used for the amplitude analyses [2, 3], in the case of $\Lambda_c^+ \rightarrow pK^-\pi^+$ decay amplitudes. We showed how the latter leads to amplitude models violating rotational invariance while the first does not. This incorrect behaviour follows from a wrong phase introduced between spin states among different decay chains.

Appendix

A. Spin in Relativistic Processes

Here, we review the description of relativistic processes involving particles with spin, which is complicated by the impossibility of a covariant definition of the spin operators [14]. Indeed, a set of spin operators \hat{S} is well defined only when acting on states describing particles at rest, so that a different set of spin operators has to be defined for each particle to express their spin states.

To describe processes involving spins of particles in relative motion, the definition of spin states must be linked to their relative kinematics. We will discuss the case of multi-body particle decays, for which the spin states of daughter particles will be referred relatively to the mother A particle states, defined in a given A rest frame reference system S_A , by specifying a sequence of Lorentz transformations.

Following Ref. [14], we consider active Lorentz transformations on the physical system: particle momenta are boosted and rotated, while the reference coordinate system (T, X, Y, Z) stays unchanged and the spatial directions specify the spin coordinate systems (the set of spin operators) of daughter particles in their rest frame. Particle momenta therefore define the spin coordinate system assigned to a given particle via the sequence of Lorentz transformations applied. To keep track of the Lorentz transformations applied, the following notation is used: a reference system S' defined in terms of another S system by applying the Lorentz transformation Λ to the momenta expressed in the S system is indicated as $S' = \Lambda S$.

The choice of the spin coordinate system for particle B moving with momentum $\mathbf{p}^A(B)$ in S_A is ambiguous, since B particle rest frames obtained by Lorentz transformations differing by rotations around $\mathbf{p}^A(B)$ represent the particle spin with different spin states. In this article, we consider the definition of the spin coordinate system for a particle at rest from that of a reference system in relative motion, while in Ref. [14], it is the spin coordinate system of the moving reference frame to be defined from that of the particle at rest. The two approaches are equivalent, but we adopt the first being more suitable for the case of particle decays, in which the initial spin coordinate system is given for the mother, rather than the daughter particles. Note that the time in the particle B frame is uniquely defined by the boost connecting different rest frames, its definition considered implicitly in the follow-

ing. There are two main choices of spin coordinate systems in literature [14]:

- (i) The canonical system S_B^C is the spin coordinate system obtained from S_A by doing a Lorentz boost $L[-\mathbf{p}^A(B)]$

$$S_B^C = L[-\mathbf{p}^A(B)]S_A. \quad (\text{A.1})$$

The canonical coordinate system is physically determined by the A momentum in the B particle rest frame $\mathbf{p}^B(A)$ requiring its direction to be opposite to that of $\mathbf{p}^A(B)$ in the A coordinate system:

$$\frac{\mathbf{p}_{x_i}^B(A)}{p^B(A)} = -\frac{\mathbf{p}_{x_i}^A(B)}{p^A(B)}. \quad (\text{A.2})$$

- (ii) The helicity system S_B^H consists in choosing the particle spin quantisation axis to be opposite to the $\mathbf{p}^B(A)$ direction. The function $\text{atan2}(y, x) \in [-\pi, \pi]$ computes the signed angle between the x -axis and the vector having components (x, y) . The helicity system for $\mathbf{p}^A(B)$ having polar and azimuthal angles θ and ϕ in the S_A system

$$\begin{aligned} \cos \theta &= \left(\frac{p_Z^A(B)}{|\mathbf{p}^A(B)|} \right), \\ \phi &= \text{atan2}(p_Y^A(B), p_X^A(B)) \end{aligned} \quad (\text{A.3})$$

is obtained by applying an $R(-\psi, -\theta, -\phi)$ Euler rotation, defined in Appendix B, aligning $\mathbf{p}^A(B)$ with the z -axis, followed by a boost $L(-p^A(B)\mathbf{Z})$:

$$S_B^H = L(-p^A(B)\mathbf{Z})R(-\psi, -\theta, -\phi)S_A. \quad (\text{A.4})$$

With this definition, the particle spin states are described in terms of the helicity states $|\mathbf{p}^A(B), s, \lambda\rangle$ introduced in Section 2. The angle ψ , associated with a rotation around $\mathbf{p}^A(B)$, determines the choice of the orthogonal spin coordinate axes. The orthogonal axes can be chosen arbitrarily but once defined must be consistently specified to avoid introducing phase differences, as shown in Section 2. Their definition is that it is better visualised as a passive rotation $R(\phi, \theta, \psi)$ applied on the (X, Y, Z) coordinate system. If there are particles other than A and B involved in the process, their momenta will provide a physical definition of the orthogonal axes, since they will change under the ψ angle rotation.

Note that any choice of the ψ angle is valid, even if the two used in the literature are $\psi = 0$ and $\psi = -\phi$. In the rest of the paper, the simplest choice $\psi = 0$ will be employed for the definition of helicity states.

A simple relation holds between helicity and canonical systems: the boost $L[-\mathbf{p}^A(B)]$ can be decomposed as

$$L[-\mathbf{p}^A(B)] = R(\phi, \theta, 0)L(-p^A(B)\mathbf{Z})R(0, -\theta, -\phi), \quad (\text{A.5})$$

so that

$$S_B^C = R(\phi, \theta, 0) S_B^H. \quad (\text{A.6})$$

The relation between canonical $|\mathbf{p}^A(B), s, \lambda\rangle$ and helicity $|\mathbf{p}^A(B), s, \lambda\rangle$ spin states, which are both defined in particle B rest frames, is given by the Wigner D -matrix (introduced in Appendix B) representing the active rotation from the helicity to the canonical system:

$$R(\phi, \theta, 0) |\mathbf{p}^A(B), s, \lambda\rangle = \sum_m D_{m,\lambda}^s(\phi, \theta, 0) |\mathbf{p}^A(B), s, m\rangle. \quad (\text{A.7})$$

Note that both canonical and helicity states are defined relatively to the given reference system S_A : canonical and helicity states defined starting from different reference systems differ by the so-called Wigner and Wick rotations, respectively [14]. When considering a chain of spin reference systems linked by canonical or helicity transformations, as in the case of multibody particle decays, the relation among spin states defined in the first and the last systems depends on the whole sequence of transformations. This fact has physical consequences, the most famous being the Thomas precession [15, 16]; in Section 4, we show how it is important to take into account the whole sequence of transformations for a correct definition of decay amplitudes.

B. Euler Rotations and Their Representation on Spin States

Here, we introduce Euler rotations and their representation on spin states, in nonrelativistic quantum mechanics, following the conventions of Ref. [17]. We consider an active rotation on a physical system described by a reference coordinate system (x, y, z) . The rotation is described by introducing a new coordinate system (x', y', z') which is rotated with respect to the reference one together with the physical system. A generic rotation of a physical system can be described by means of Euler rotations, which, in the $z-y-z$ convention for the rotation axes, are defined as

$$R(\alpha, \beta, \gamma) = R_z(\alpha) R_y(\beta) R_z(\gamma) = e^{-i\alpha S_{Az}} e^{-i\beta S_{Ay}} e^{-i\gamma S_{Az}}, \quad (\text{B.1})$$

in which rotations are expressed with respect to the original axes (x, y, z) . The three Euler angles α, β, γ can be computed from the unit vectors describing the rotated system in the terms of the original coordinates as

$$\alpha = \text{atan2}\left(z'_y, z'_x\right) \in [-\pi, \pi], \quad (\text{B.2})$$

$$\beta = \arccos\left(z'_z\right) \in [0, \pi], \quad (\text{B.3})$$

$$\gamma = \text{atan2}\left(y'_z, -x'_z\right) \in [-\pi, \pi]. \quad (\text{B.4})$$

The action of a Euler rotation on spin states $|s, m\rangle$ associated with the spin coordinate system (x, y, z) is

$$\hat{R}(\alpha, \beta, \gamma) |s, m\rangle = \sum_{m'=-s}^s D_{m',m}^s(\alpha, \beta, \gamma) |s, m'\rangle, \quad (\text{B.5})$$

in which the Wigner D -matrices $D_{m',m}^s(\alpha, \beta, \gamma)$ are

$$D_{m',m}^s(\alpha, \beta, \gamma) = \langle s, m' | R(\alpha, \beta, \gamma) |s, m\rangle. \quad (\text{B.6})$$

The Wigner D -matrices of index s are spin s representations of the $SU(2)$ group. Following Equation (B.1), the Wigner D -matrices can be factorised as

$$\begin{aligned} D_{m',m}^s(\alpha, \beta, \gamma) &= \langle s, m' | e^{-i\alpha S_{Az}} e^{-i\beta S_{Ay}} e^{-i\gamma S_{Az}} |s, m\rangle \\ &= e^{-im'\alpha} d_{m',m}^s(\beta) e^{-im'\gamma}, \end{aligned} \quad (\text{B.7})$$

in which the Wigner D -matrix elements are real combinations of trigonometric functions of β and their analytical expression is reported in Ref. [17]. Wigner D -matrices have many properties following from those of the rotation group; for the purpose of this article, we report

$$D_{m',m}^s(\alpha, \beta, \gamma) = D_{m,m'}^s(\gamma, -\beta, \alpha), \quad (\text{B.8})$$

$$D_{m',m}^{*s}(\alpha, \beta, \gamma) = D_{m',m}^s(-\alpha, \beta, -\gamma). \quad (\text{B.9})$$

The inverse rotation from the final coordinate system (x', y', z') to the initial one (x, y, z) follows from Equation (B.1):

$$\begin{aligned} R^{-1}(\alpha, \beta, \gamma) &= R_z^{-1}(\gamma) R_y^{-1}(\beta) R_z^{-1}(\alpha) = e^{i\gamma S_{Az}} e^{i\beta S_{Ay}} e^{i\alpha S_{Az}} \\ &= R(-\gamma, -\beta, -\alpha), \end{aligned} \quad (\text{B.10})$$

and the Wigner D -matrix representation on $|s, m\rangle$ states is $D_{m',m}^s(-\gamma, -\beta, -\alpha)$. This definition of inverse Euler rotation allows undoing step by step the three rotations comprising the Euler rotation. The usual inverse Euler rotation, given in terms of the positive β angle, inverts the rotation following a different path, which can introduce additional 2π rotations leading to inequivalent trajectories in the $SU(2)$ group representing spin. This issue is relevant for fermion states, as mentioned in Section 2.

As a practical example, let us consider the inverse of a rotation of angle θ around the y -axis, $R(0, \theta, 0)$, which can be chosen to be $R(0, -\theta, 0)$ or $R(\pi, \theta, \pi)$. The Wigner D -matrix representing $\hat{R}(0, -\theta, 0)$ on a spin 1/2 state is

$$D_{m',m}^{1/2}(0, -\theta, 0) = d_{m',m}^{1/2}(-\theta) = \begin{pmatrix} \cos \frac{\theta}{2} & \sin \frac{\theta}{2} \\ -\sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix}, \quad (\text{B.11})$$

while that representing $\widehat{R}(\pi, \theta, \pi)$ is

$$D_{m',m}^{1/2}(\pi, \theta, \pi) = e^{-i\pi(m+m')} d_{m',m}^{1/2}(-\theta) \\ = \begin{pmatrix} -\cos \frac{\theta}{2} & -\sin \frac{\theta}{2} \\ \sin \frac{\theta}{2} & -\cos \frac{\theta}{2} \end{pmatrix} = -D_{m',m}^{1/2}(0, \theta, 0), \quad (\text{B.12})$$

which has the opposite sign with respect to the first matrix.

C. Polarised Differential Decay Rate

The generic spin state of a statistical ensemble of particles is defined by the associated density operator $\widehat{\rho}$ [14]: given an ensemble of spin states $|\psi\rangle_i$ occurring with probability p_i , the density operator is

$$\widehat{\rho} = \sum_i p_i |\psi\rangle_i \langle \psi|_i, \quad (\text{C.1})$$

so that the expectation value of any operator \widehat{X} can be expressed as

$$\langle \widehat{X} \rangle = \sum_i p_i \langle \psi | \widehat{X} | \psi \rangle_i = \text{Tr}[\widehat{\rho} \widehat{X}]. \quad (\text{C.2})$$

From the definition, Equation (C.1), follows that the density operator is Hermitian with a unit trace.

The decay rate of multibody decays $A \rightarrow \{i = 1, \dots, n\}$ for definite spin eigenstates is the squared modulus of the transition amplitude between the A particle initial state $|s_A, m_A\rangle$ and the final particle product state $|\{s_i\}, \{m_i\}\rangle = \otimes_i |s_i, m_i\rangle$:

$$p_{m_A, \{m_i\}}(\Omega) = |\langle s_A, m_A | T \wedge | \{s_i\}, \{m_i\} \rangle|^2 = \left| \mathcal{A}_{m_A, \{m_i\}}(\Omega) \right|^2. \quad (\text{C.3})$$

Generic polarisation states are described by introducing the density operators for the initial particle state $\rho^{\wedge A}$ and the final particle states $\rho^{\wedge \{i\}}$, which are included in the decay rate (Equation (C.3)) by inserting suitable identity resolutions, obtaining

$$p(\rho^{\wedge A}, \rho^{\wedge \{i\}}; \Omega) = \text{tr} \left[\rho^{\wedge A} \widehat{T} \rho^{\wedge \{i\}} T \wedge^\dagger \right] \\ = \sum_{m_A, m'_A, \{m_i\}, \{m'_i\}} \widehat{\rho}_{m_A, m'_A}^A \widehat{\rho}_{\{m_i\}, \{m'_i\}}^{\{i\}} \mathcal{A}_{m_A, \{m_i\}} \\ \cdot (\Omega) \mathcal{A}_{m'_A, \{m'_i\}}^*(\Omega). \quad (\text{C.4})$$

D. $\Lambda_c^+ \rightarrow pK^- \pi^+$ Amplitude Model

The $\Lambda_c^+ \rightarrow pK^- \pi^+$ amplitude model is built closely following the three-body helicity amplitudes described in Section 4,

TABLE 1: Helicity couplings and Breit-Wigner parameter values employed in the numerical study in Section 5.

Parameter	Value
K^*	
$H_{1/2,0}$	1
$H_{1/2,-1}$	$0.5 + 0.5i$
$H_{-1/2,1}$	i
$H_{-1/2,0}$	$-0.5 - 0.5i$
$m(\text{GeV})$	0.9
$\Gamma(\text{GeV})$	0.2
Λ^*	
$H_{-1/2,0}$	i
$H_{1/2,0}$	$0.8 - 0.4i$
$m(\text{GeV})$	1.6
$\Gamma(\text{GeV})$	0.2
Δ^*	
$H_{-1/2,0}$	$0.6 - 0.4i$
$H_{1/2,0}$	$0.1i$
$m(\text{GeV})$	1.4
$\Gamma(\text{GeV})$	0.2

with the identifications $A \leftrightarrow \Lambda_c^+$, $1 \leftrightarrow p$, $2 \leftrightarrow K^-$, $3 \leftrightarrow \pi^+$, $R \leftrightarrow \Lambda^*$, $S \leftrightarrow \Delta^{*++}$, and $U \leftrightarrow K^*$. In this appendix, we report the definition of the helicity amplitudes for the specific $\Lambda_c^+ \rightarrow pK^- \pi^+$ decay case.

Starting from the decay chain $\Lambda_c^+ \rightarrow pK^* (\rightarrow K^- \pi^+)$, the weak decay $\Lambda_c^+ \rightarrow pK^*$ is described by

$$\mathcal{A}_{m_{\Lambda_c^+}, \lambda_p, \bar{\lambda}_{K^*}}^{\Lambda_c^+ \rightarrow pK^*} = \mathcal{H}_{\lambda_p, \bar{\lambda}_{K^*}}^{\Lambda_c^+ \rightarrow pK^*} D_{m_{\Lambda_c^+}, \lambda_p, \bar{\lambda}_{K^*}}^{*1/2}(\phi_p, \theta_p, 0), \quad (\text{D.1})$$

in which proton and K^* helicities λ_p and $\bar{\lambda}_{K^*}$ are defined in the proton helicity frame reached from the Λ_c^+ baryon polarisation frame. For spin zero K^* resonances, the angular momentum conservation relations (Equation (17)) allow two complex couplings corresponding to $m_p = \pm 1/2$; for higher spin resonances, four couplings are allowed, corresponding to $\{m_p = 1/2; \bar{\lambda}_{K^*} = 0, -1\}$ and $\{m_p = -(1/2); \bar{\lambda}_{K^*} = 0, 1\}$. The couplings are independent of each other because of parity violation in weak decays. The strong decay $K^* \rightarrow K^- \pi^+$ contribution is

$$\mathcal{A}_{\bar{\lambda}_{K^*}}^{K^* \rightarrow K^- \pi^+} = \mathcal{H}_{0,0}^{K^* \rightarrow K^- \pi^+} D_{\bar{\lambda}_{K^*}, 0}^{*J_{K^*}}(\bar{\phi}_K, \bar{\theta}_K, 0) \mathcal{R}(m_{K^- \pi^+}^2), \quad (\text{D.2})$$

in which $\bar{\phi}_K$ and $\bar{\theta}_K$ are the kaon azimuthal and polar angles in the K^* opposite-helicity frame (obtained boosting to the K^* rest frame after the rotation $R(\phi_p, \theta_p, 0)$). The function $\mathcal{R}(m_{K^- \pi^+}^2)$ describes the nonnegligible mass width of the

intermediate state. The coupling $\mathcal{H}_{0,0}^{K^* \rightarrow K^- \pi^+}$ can not be determined independently of $\mathcal{H}_{m_p, \lambda_{K^*}}^{\Lambda_c^+ \rightarrow K^* p}$ couplings; thus, it is set equal to 1 and absorbed into the latter.

Considering the decay chain $\Lambda_c^+ \rightarrow \Lambda^* (\rightarrow p K^-) \pi^+$, the weak decay $\Lambda_c^+ \rightarrow \Lambda^* \pi^+$ is described by

$$\mathcal{A}_{m_{\Lambda_c^+}, \lambda_{\Lambda^*}}^{\Lambda_c^+ \rightarrow \Lambda^* \pi^+} = \mathcal{H}_{\lambda_{\Lambda^*}, 0}^{\Lambda_c^+ \rightarrow \Lambda^* \pi^+} D_{m_{\Lambda_c^+}, \lambda_{\Lambda^*}}^{*1/2}(\phi_{\Lambda^*}, \theta_{\Lambda^*}, 0), \quad (\text{D.3})$$

with λ_{Λ^*} defined in the Λ^* helicity system reached from the Λ_c^+ reference frame. The angular momentum conservation relations (Equation (17)) allow two helicity couplings, $\lambda_{\Lambda^*} = \pm 1/2$, to fit for each resonance whatever J_{Λ^*} is. The strong decay $\Lambda_c^+ \rightarrow p K^-$ is described by

$$\mathcal{A}_{\lambda_{\Lambda^*}, \lambda_p^{\Lambda^*}}^{\Lambda_c^+ \rightarrow p K^-} = \mathcal{H}_{\lambda_p^{\Lambda^*}, 0}^{\Lambda_c^+ \rightarrow p K^-} D_{\lambda_{\Lambda^*}, \lambda_p^{\Lambda^*}}^{*J_{\Lambda^*}}(\phi_p^{\Lambda^*}, \theta_p^{\Lambda^*}, 0) \mathcal{R}(m_{pK^-}^2), \quad (\text{D.4})$$

with $\lambda_p^{\Lambda^*}$ as the proton helicity in the system reached from the Λ^* helicity frame. Parity conservation in strong decays relates the two helicity couplings corresponding to $\lambda_p^{\Lambda^*} = \pm 1/2$ by

$$\mathcal{H}_{-\lambda_p^{\Lambda^*}, 0}^{\Lambda_c^+ \rightarrow p K^-} = -P_{\Lambda^*} (-1)^{J_{\Lambda^*} - 1/2} \mathcal{H}_{\lambda_p^{\Lambda^*}, 0}^{\Lambda_c^+ \rightarrow p K^-}, \quad (\text{D.5})$$

in which P_{Λ^*} is the parity of the Λ^* resonance and the proton and kaon parities $P_p = 1$, $P_K = -1$ have been inserted. They are absorbed into $\mathcal{H}_{\lambda_{\Lambda^*}, 0}^{\Lambda_c^+ \rightarrow \Lambda^* \pi^+}$ couplings by setting

$$\mathcal{H}_{+1/2, 0}^{\Lambda_c^+ \rightarrow p K^-} = 1, \quad \mathcal{H}_{-1/2, 0}^{\Lambda_c^+ \rightarrow p K^-} = -P_{\Lambda^*} (-1)^{J_{\Lambda^*} - 1/2}. \quad (\text{D.6})$$

Considering the third decay chain $\Lambda_c^+ \rightarrow \Delta^{++} (\rightarrow p \pi^+) K$, the weak decay $\Lambda_c^+ \rightarrow \Delta^{++} K^-$ is described by

$$\mathcal{A}_{m_{\Lambda_c^+}, \lambda_{\Delta^*}}^{\Lambda_c^+ \rightarrow \Delta^* K^-} = \mathcal{H}_{\lambda_{\Delta^*}, 0}^{\Lambda_c^+ \rightarrow \Delta^* K^-} D_{m_{\Lambda_c^+}, \lambda_{\Delta^*}}^{*1/2}(\phi_{\Delta^*}, \theta_{\Delta^*}, 0), \quad (\text{D.7})$$

with λ_{Δ^*} defined in the Δ^* helicity system reached from the Λ_c^+ frame. The strong decay $\Delta^{++} \rightarrow p \pi^+$ amplitude is written as

$$\mathcal{A}_{\lambda_{\Delta^*}, \lambda_p^{\Delta^*}}^{\Delta^{++} \rightarrow p \pi^+} = \mathcal{H}_{\lambda_p^{\Delta^*}, 0}^{\Delta^{++} \rightarrow p \pi^+} D_{\lambda_{\Delta^*}, \lambda_p^{\Delta^*}}^{*J_{\Delta^*}}(\phi_p^{\Delta^*}, \theta_p^{\Delta^*}, 0) \mathcal{R}(m_{p\pi^+}^2), \quad (\text{D.8})$$

with $\lambda_p^{\Delta^*}$ defined in the proton helicity system reached from the Δ^* helicity frame. Helicity couplings are defined as for the Λ^* decay chain.

The decay amplitudes for each decay chain are obtained as in Equations (22) and (23), to which the rotations needed to match the proton spin state definitions must be applied, as in Equations (45) and (46).

For the numerical study presented in Section 5, we consider a $\Lambda_c^+ \rightarrow p K^- \pi^+$ amplitude model consisting of three resonances, one per decay channel, with invariant mass dependence described by relativistic Breit-Wigner functions, characterised by mass and width parameters. We consider the following spin-parity J^P assignments: $K^*(1^-)$, $\Lambda^*(1/2^-)$,

and $\Delta^*(1/2^-)$. The values of the complex couplings and Breit-Wigner mass and width parameters employed in the study are reported in Table 1: they are chosen in order to produce significant interference effects.

Data Availability

The articles used to support the findings of this study are included within the article and are cited at relevant places within the text as references.

Conflicts of Interest

The author declares that there are no conflicts of interest.

Acknowledgments

I thank Mikhail Mikhaseenko and Alessandro Pilloni for the critical review of the first version of the article, in particular for pointing out the need of a Wigner rotation in the matching of the final particle spin states. I thank my colleagues Louis Henry, Fernando Martínez Vidal, Andrea Merli, Nicola Neri, and Elisabetta Spadaro Norella for their interesting discussions about the helicity formalism and its application to baryon decays. This work was supported by the ERC Consolidator Grant SELDOM G.A. 771642.

References

- [1] M. Jacob and G. C. Wick, "On the general theory of collisions for particles with spin," *Annals of Physics*, vol. 7, no. 4, pp. 404–428, 1959.
- [2] R. Aaij, B. Adeva, M. Adinolfi et al., "Observation of $J/\psi p$ resonances consistent with pentaquark states in $\Lambda_b^0 \rightarrow J/\psi K^- p$ decays," *Physical Review Letters*, vol. 115, no. 7, article 072001, 2015.
- [3] R. Mizuk, R. Chistov, I. Adachi et al., "Observation of two resonancelike structures in the $\pi^+ \chi_{c1}$ mass distribution in exclusive $\bar{B}^0 \rightarrow K^- \chi_{c1}$ decays," *Physical Review D*, vol. 78, no. 7, article 072004, 2008.
- [4] H. Chen and R.-G. Ping, "Coherent helicity amplitude for sequential decays," *Physical Review D*, vol. 95, no. 7, article 076010, 2017.
- [5] M. Mikhaseenko, M. Albaladejo, Ł. Bibrzycki et al., "Dalitz-plot decomposition for three-body decays," *Physical Review D*, vol. 101, no. 3, article 034033, 2020.
- [6] J. J. Sakurai, *Modern Quantum Mechanics*, Addison-Wesley Publishing Company, 1994.
- [7] E. Gourghoulon, *Special Relativity in General Frames: From Particles to Astrophysics*, Springer, Berlin, Heidelberg, 2013.
- [8] H. Rauch, A. Zeilinger, G. Badurek, A. Wilfling, W. Bauspiess, and U. Bonse, "Verification of coherent spinor rotation of fermions," *Physics Letters A*, vol. 54, no. 6, pp. 425–427, 1975.
- [9] S. A. Werner, R. Colella, A. W. Overhauser, and C. F. Eagen, "Observation of the phase shift of a neutron due to precession in a magnetic field," *Physical Review Letters*, vol. 35, no. 16, pp. 1053–1055, 1975.
- [10] "TensorFlowAnalysis: a collection of useful functions and example scripts for performing amplitude fits using TensorFlow," <https://gitlab.cern.ch/poluekt/TensorFlowAnalysis>.

- [11] D. Marangotto, *Amplitude analysis and polarisation measurement Λ_c^+ of the baryon $pK^- \pi^+$ in final state for electromagnetic dipole moment experiment*, Università degli studi di Milano, 2020, <https://cds.cern.ch/record/2713231>.
- [12] M. Abadi, A. Agarwal, P. Barham et al., “TensorFlow: large-scale machine learning on heterogeneous systems,” 2015, <https://www.tensorflow.org>.
- [13] R. Brun and F. Rademakers, “ROOT: an object oriented data analysis framework,” *Nuclear Instruments and Methods in Physics Research*, vol. A389, pp. 81–86, 1997.
- [14] E. Leader, “Spin in particle physics,” *Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology*, vol. 15, 2011.
- [15] L. H. Thomas, “The motion of the epinning electron,” *Nature*, vol. 117, no. 2945, p. 514, 1926.
- [16] L. H. Thomas, “I. The kinematics of an electron with an axis,” *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, vol. 3, no. 13, pp. 1–22, 2009.
- [17] J. D. Richman, “An experimenter's guide to the helicity formalism,” 1984.