

## Research Article

# Neutrophil Secretion Induced by an Intracellular $\text{Ca}^{2+}$ Rise and Followed by Whole-Cell Patch-Clamp Recordings Occurs Without any Selective Mobilization of Different Granule Populations

Daniel Granfeldt,<sup>1,2</sup> Olle Harbecke,<sup>1,3</sup> Åse Björstad,<sup>1</sup> Anna Karlsson,<sup>1</sup> and Claes Dahlgren<sup>1</sup>

<sup>1</sup>The Phagocyte Research Laboratory, Department of Rheumatology and Inflammation Research, Göteborg University, Guldhedsgatan 10, 413 46 Göteborg, Sweden

<sup>2</sup>Department of Chemistry & Bioscience, Chalmers University of Technology, Kemivägen 10, 412 96 Göteborg, Sweden

<sup>3</sup>Univa AB, Lunds Universitet, 221 00 Lund, Sweden

Received 20 December 2005; Revised 22 March 2006; Accepted 4 May 2006

We have investigated calcium-induced secretion in human neutrophils, using a whole-cell patch-clamp technique. Mobilization of subcellular granules to the cell membrane was followed as the change in membrane capacitance ( $\Delta C_m$ ). Both the magnitude and the kinetics of the response differed between low and high concentrations of  $\text{Ca}^{2+}$ . A sustained secretion following a short lag phase was induced by high concentrations of  $\text{Ca}^{2+}$  (100  $\mu\text{M}$  and higher). A stable plateau was reached after 5–7 minutes at  $\Delta C_m$  values corresponding to values expected after all specific as well as azurophil granules have been mobilized. Capacitance values of the same magnitude could be obtained also at lower  $\text{Ca}^{2+}$  concentrations, but typically no stable plateau was reached within the measuring time. In contrast to previous studies, we were unable to detect any pattern of secretion corresponding to a distinct submaximal response or selective mobilization of granule subsets specified by their  $\text{Ca}^{2+}$ -sensitivity.

Copyright © 2006 Daniel Granfeldt et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Neutrophils constitute one of the major cell types of our innate immune system, with a main task to phagocytose and kill invading microbial pathogens [1]. To perform their duties, the neutrophils are equipped with subcellular granules, intracellular storage vesicles that contain their destructive weaponry (ie, microbicidal and proinflammatory agents), and a reserve pool of a variety of cell-surface receptors [2]. Neutrophils contain four granule/vesicle types, the azurophil granules, the specific granules, the gelatinase granules, and the secretory vesicles. These are formed (in the order listed) during neutrophil maturation in the bone marrow [3]. During neutrophil activation and diapedesis, the content of these granule/vesicle populations is secreted in a defined, hierarchical manner, in the reverse order of their formation [4].

A common model for membrane fusion events has been suggested [5, 6], and homologues to several (but not all) proteins of the synaptic secretory machinery have been found also in neutrophils [7–9]. Even though the precise molecular

regulatory mechanism of exocytosis might differ between cell types, a common feature is the activation of the secretory system by an elevation of the cytosolic concentration of free calcium ions  $[\text{Ca}^{2+}]_i$  [10], and the role of  $\text{Ca}^{2+}$  as a regulator of granule mobilization in neutrophils has been recognized for several years [11]. The regulatory system in neutrophils is particularly interesting, as these cells contain several different granule/vesicle populations that should be released sequentially [3]. There are several reports suggesting that the distinct neutrophil granule populations have different  $\text{Ca}^{2+}$  requirements for fusion with the plasma membrane [12–14]. Mobilization of the azurophil granule population bears resemblance to neuronal exocytosis in that it is characterized by low affinity for  $\text{Ca}^{2+}$  and a high degree of cooperativity. In contrast, fusion with the plasma membrane of the more easily mobilized granules was suggested to resemble that of endocrine exocytosis in that it is characterized by high affinity for  $\text{Ca}^{2+}$  and low cooperativity [15]. Based on results obtained from patch-clamp-capacitance studies a difference in calcium affinity among neutrophil granule population has

also been described [16]. This is a unique feature for the neutrophil granules and suggested to be the basis for  $\text{Ca}^{2+}$  induced selective granule secretion in neutrophils [15].

Neutrophil secretion is tightly regulated by the cytoskeleton [17] and it should be noticed that the conclusion that neutrophils are unique in the sense that they contain granule populations displaying distinct differences with respect to their  $\text{Ca}^{2+}$ -affinities was drawn from experiments in which cytoskeleton disrupting agents (cytochalasin B) were included. An intact cytoskeleton will most likely have profound impact on the process of granule release and these conclusions therefore need to be reevaluated. In this study, using whole-cell capacitance measurements of secretion in neutrophils with intact cytoskeletons, we found no evidence for the coexistence of two separate control mechanisms for granule mobilization.

## METHODS

### Isolation of phagocytic cells

Neutrophils were isolated [18] from heparinized whole blood obtained from apparently healthy adults. After dextran sedimentation, hypotonic lysis of the remaining erythrocytes, and centrifugation in a Ficoll-Paque gradient, the neutrophils were washed and resuspended in Krebs-Ringer phosphate buffer containing glucose (10 mM),  $\text{Ca}^{2+}$  (1 mM), and  $\text{Mg}^{2+}$  (1.5 mM) (KRG, pH 7.3). The cells were kept on melting ice and used within six hours.

### Whole-cell patch-clamp capacitance measurement of secretion

Patch-clamp recordings were performed according to standard procedures [19]. Coverslips were incubated with a bovine serum albumin (BSA) solution (1% w/v in distilled water) for 10 min and then washed with the standard bath solution (BS; see below) before use. Neutrophils were allowed to adhere to the slides for 5–10 min, and nonadherent cells were washed away with BS. Patch pipettes had pipette resistances between 4–7 M $\Omega$ . Capacitance changes were recorded using an EPC-9 patch-clamp amplifier (HEKA, Lambrecht/Pfalz, Germany). From a holding potential of  $-10$  mV, the cells were exposed to a sine-shaped, alternating voltage pulse [20] with a frequency of 1.0 kHz and amplitude of 20 mV. From the resulting current, the capacitance was determined by the lock-in software-based phase-sensitive detector [21, 22] and sampled by the X-chart extension of pulse 8.41 (HEKA). The membrane capacitance was registered for 10 min or (if possible) longer. Only recordings during which the access resistance was well below 20 M $\Omega$  when the whole cell configuration was established, as well as during the major part of the recording, were regarded as reliable. Baseline or background secretion was determined using a pipette solution containing no added  $\text{Ca}^{2+}$  and 2 mM of the  $\text{Ca}^{2+}$  chelator HEDTA. This solution was regarded as  $\text{Ca}^{2+}$  free. The capacitance recordings were started when the membrane was ruptured and the whole cell configuration was established.

The initial value (typically around 2.5 pF) of the capacitance for each cell has been subtracted in the recordings shown.

### Data presentation and analysis

The time of membrane rupture and establishment of the whole cell configuration was set as time zero and initial capacitance at this time was about 2.5 pF ( $2.57 \pm 0.05$  pF, mean  $\pm$  SEM,  $n = 140$ ). X-chart data were imported into IGOR 3.03 for Macintosh (Wave Metrics Inc, Lake Oswego, OR) and analyzed. Statistical calculations were performed using Prism 3.02 (Graph Pad Software Inc, San Diego, CA) and  $P < .05$  was considered significant.

### Solutions and reagents

The standard bath solution (BS) contained 140 mM NaCl, 5 mM KCl, 1 mM  $\text{MgCl}_2$ , 2 mM  $\text{CaCl}_2$ , 10 mM glucose, and 10 mM HEPES, pH 7.4. The standard pipette solution (PS) contained 112 mM K-glutamate, 18 mM NaCl, 1.8 mM  $\text{MgCl}_2$ , 2.5 mM  $\text{Na}_2\text{ATP}$ , and 9 mM HEPES, pH 7.2. The pipette solution was supplemented with suitable concentrations of calcium ions and calcium chelator (citrate or HEDTA) to reach the desired concentration of free calcium, as determined by WinMaxChelator 1.78 (Chris Patton, Stanford, CA, USA), using the BERS constants [23]. Dextran and Ficoll-Paque were purchased from Pharmacia (Uppsala, Sweden).

## RESULTS

Neutrophils respond to increased levels of  $[\text{Ca}^{2+}]_i$  with exocytosis of granules (degranulation), in this system manifested as an increased membrane capacitance. The degree of granule mobilization depends on the concentration of free  $\text{Ca}^{2+}$  used in the pipette solution, which in the study ranged from nominally calcium-free solutions to 300  $\mu\text{M}$ . The maximal  $\Delta C_m$  values obtained at high  $[\text{Ca}^{2+}]_i$  were 6–7 pF (Figure 1(a)), which correspond to an increase in total membrane area approaching 300%. The magnitude of the capacitance change as well as the kinetics of the secretory response differed between the systems with a low (10  $\mu\text{M}$  or lower) or a high (100  $\mu\text{M}$  or higher) concentration of  $\text{Ca}^{2+}$  in the pipette, (Figures 1(a) and 1(b), Table 1).

The secretory response induced by a high  $[\text{Ca}^{2+}]_i$  started with a lag phase during which the capacitance increase was around 10 fF/sec (Figure 1(b)). The lag phase was followed by a more sustained phase of secretion after which the capacitance recordings reached a plateau, usually 5–7 min after rupture of the membrane. The maximal obtainable rate of secretion using a lower  $[\text{Ca}^{2+}]_i$ , (0, 1, and 10  $\mu\text{M}$ ) was around 10 fF/s, whereas this rate was around 50–60 fF/s for the higher concentrations (100 and 300  $\mu\text{M}$ ). No difference in maximal rate of secretion was found between  $[\text{Ca}^{2+}]_i$  of 100 and 300  $\mu\text{M}$ , respectively (Figure 2(a)). However, the time required to reach a capacitance value corresponding to half of the maximal (plateau) value was significantly ( $P < .001$ ) shortened when the  $[\text{Ca}^{2+}]_i$  concentration was increased (Figure 2(b)). The initial rate of secretion was higher with




FIGURE 1: Characteristics of calcium-induced secretion in neutrophils. (a) Change in membrane capacitance induced by increasing concentrations of  $[Ca^{2+}]_i$ . Values were determined when a plateau level was reached or 10 min after breakthrough with the patch pipette. (b) Time-course of calcium induced neutrophil exocytosis. The recordings are from representative experiments.

TABLE 1: Basic characteristics of the secretory process in human neutrophils.

$[Ca^{2+}]_i$ ( $\mu M$ )	$n$	$C_{breakthrough}$ (pF) <sup>a</sup>	$\Delta C_m$ (pF) <sup>b</sup>	Rate <sub>max</sub> (fF/s) <sup>c</sup>	Point of inflection (pF) <sup>c</sup>	Half-time rise (s) <sup>d</sup>
0	13	$2.42 \pm 0.36$	$1.10 \pm 0.44$	$7.38 \pm 2.69$	nd <sup>e</sup>	nd
1	7	$2.20 \pm 0.24$	$1.39 \pm 0.48$	$6.63 \pm 1.86$	nd	nd
10	17	$2.40 \pm 0.35$	$1.82 \pm 0.59$	$5.27 \pm 3.06$	nd	nd
100	23	$2.62 \pm 0.50$	$5.95 \pm 1.35$	$51.55 \pm 18.52$	$0.57 \pm 0.32$	$142.2 \pm 52.0$
300	27	$2.71 \pm 0.73$	$6.30 \pm 1.29$	$58.90 \pm 22.73$	$0.35 \pm 0.17$	$95.8 \pm 31.4$

Secretion was induced by an introduction through the patch pipette of pipette solutions containing increased concentrations of free  $Ca^{2+}$ . The concentration of  $Ca^{2+}$  was varied from 0 to 300  $\mu M$  and all numerals presented represent mean values  $\pm$  S.D. The number of cells studied ( $n$ ) is given for each concentration of  $Ca^{2+}$ .

<sup>a</sup>Values represent the initial capacitance values of the cell, determined upon the establishment of the whole-cell configuration.

<sup>b</sup>Values represent the change in capacitance reached at the plateau (for calcium concentrations of 100 and 300  $\mu M$ ), and for comparison with those that did not reach a plateau (calcium concentrations 1 and 10  $\mu M$ ) 5 minutes after breakthrough, a time point that corresponds to the time required to reach a plateau at the higher calcium concentrations.

<sup>c</sup>Values represent the maximal rate of capacitance change and the capacitance value (point of inflection) corresponding to the time point at which the maximal of secretion was reached, respectively.

<sup>d</sup>Values represent the time required to reach a  $\Delta C_m$  corresponding to 50% of the plateau value.

<sup>e</sup>Not determined.

a  $[Ca^{2+}]_i$  of 300  $\mu M$  compared to 100  $\mu M$  and the time required to reach a maximal response was shorter. The  $\Delta C_m$  value at which the maximal rate of secretion reached was somewhat lower for the responses induced at a  $[Ca^{2+}]_i$  of 300  $\mu M$  as compared to 100  $\mu M$ . These values were typically below 0.5 pF and almost always (48 of 50 cells, 96%) below 1.0 pF for both concentrations (Table 1, point of inflection).

When the patch pipette contained lower  $Ca^{2+}$  concentrations (up to 10  $\mu M$ ), no distinct shift in the rate of secretion could be observed (Figure 1(b)). The secretory response induced at lower  $[Ca^{2+}]_i$  also differed from that induced with higher  $[Ca^{2+}]_i$  in that it seldom reached a plateau during the 10 min measurement time. In fact, four individual measurements that maintained the required values of seal and access resistance for around 15 minutes ( $875 \pm 171$ s,  $n = 4$ ) still showed increasing capacitance that could reach  $\Delta C_m$  values of up to 5 pF (mean =  $4.27 \pm 0.85$  pF,  $n = 4$ ) without any

sign of decrease in the secretion rate (Figure 3). Perfusion with nominally calcium-free solutions resulted in an increase in capacitance during a 3–5 min period, and this phase typically reached a  $\Delta C_m$  value of around 1 pF and was followed by a stable and prolonged plateau level. The outcome of experiments with a  $[Ca^{2+}]_i$  of 60  $\mu M$  was highly unpredictable. Such an experiment could result in either of the two secretory patterns described (high or low  $[Ca^{2+}]_i$ ), suggesting that there is a threshold concentration somewhere around 60  $\mu M$ .

Obtaining the rate of capacitance increase directly by plotting the derivate of the capacitance traces against time revealed a unimodal pattern of response with only one distinct peak, for both concentrations (Figure 4). Furthermore, all the traces from both 100  $\mu M$  and 300  $\mu M$   $[Ca^{2+}]_i$  could be well fitted ( $R^2 = 0.996 \pm 0.001$ , mean  $\pm$  SEM,  $n = 50$ ) to a simple logistic equation describing a monophasic curve with sigmoid appearance.




FIGURE 2: Dynamics of secretion at different  $[Ca^{2+}]_i$ . (a) Maximal rates of capacitance increase during neutrophil exocytosis. The capacitance values were recorded during a 5–10 min period after breakthrough with the patch pipettes, and the maximal rate of exocytosis was determined from each capacitance trace. (b) Higher  $[Ca^{2+}]_i$  gives shorter half-rise time. The time required for reaching half of the maximal capacitance increase is induced by 100  $\mu M$  and 300  $\mu M$   $[Ca^{2+}]_i$ , respectively.




FIGURE 3: Long exposure to low levels of  $[Ca^{2+}]_i$  can induce secretion of all granules. The change in membrane capacitance induced with a pipette solution containing 10  $\mu M$  of  $[Ca^{2+}]_i$ . The curve is from one of the four recordings that could be followed for more than 15 min.

## DISCUSSION

The level of cytosolic  $Ca^{2+}$  is a key regulatory element for the release of neutrophil granule markers and exposure of new membrane proteins [11]. The hierarchy in neutrophil vesicle and granule mobilization has been well established in vitro as well as in vivo [4], and as a general rule the degree of mobilization is inversely proportional to the size of the granules. A fairly simple model for how this hierarchy is achieved has been presented [24]. According to this model the densest (and largest in size) granule population (the azurophil granules) is also the hardest to mobilize because a greater

stimulation or pulling force is required for their fusion with the plasma membrane. Such a pulling force can be manifested either in the form of highly elevated  $Ca^{2+}$  concentrations or prolonged exposure to lower levels. We achieve a  $\Delta C_m$  of 6–7 pF through microperfusion of neutrophils with  $[Ca^{2+}]_i$  in the 100–300  $\mu M$  range corresponding to earlier published values for complete degranulation [15, 25] where specific granules are suggested to account for 2.4 pF and azurophil granules for 3.6 pF, respectively [26].

We have chosen to focus on the two secretion patterns obtained with high ( $>100 \mu M$ ) and low (10  $\mu M$ )  $[Ca^{2+}]_i$ . Results obtained from measurements with intermediate calcium concentrations also show one or another of the two secretion patterns. This variability probably depends on whether or not the specific cell experiences a concentration of calcium that is high enough to induce a rapid secretory response, but more importantly, the high degree of mobilization was never associated with a biphasic pattern of capacitance change as previously reported [15], as determined by the derivation of the capacitance traces. The high- and low- $Ca^{2+}$ -affinity model proposed to be the regulatory base for the hierarchical neutrophil granule secretion [15] implies that the release of azurophil granules occurring at a high rate ( $>50$ – $60$  fF/s) cannot be achieved until all peroxidase negative granules have been mobilized. This suggests that high rates of secretion should not occur until  $\Delta C_m$  exceeds 2.4 pF, a value corresponding to the membrane area of all the specific granules. We found, however, that a high rate ( $>70$  fF/s) of secretion could be obtained long before  $\Delta C_m$  had reached 2.4 pF. In fact, the point of inflection was almost always (48 out of 50 cells, 96%) found to be below 1 pF and could even be as low as 0.1 pF. Our data thus suggest that also the specific granules can be secreted at a high rate.

Furthermore, our data also imply that the rate of secretion is not directly linked to the type of granule involved




FIGURE 4: Neutrophil secretion is monophasic. The rate of capacitance increase during neutrophil exocytosis is induced by  $[Ca^{2+}]_i$  of 100  $\mu M$  (left) and 300  $\mu M$  (right), respectively. (a) and (b) show the capacitance recordings of two representative experiments, and (c) and (d) show the rates of exocytosis, determined as the derivatives of these capacitance curves. No cells ( $n = 50$ ) showed two peaks in the derived traces. No direct comparisons were made between the two pairs of traces (a and c versus b and d) making it possible to use scales that accurately show each individual trace.

in the secretory process. This suggestion gains support from the fact that  $\Delta C_m$  values close to those corresponding to a mobilization of all granules (including at least 50% of the azurophil granules) can be obtained with an experimental setup in which the rate of secretion is fairly low ( $< 10$  fF/s). Based on these data it also seems reasonable to believe that the azurophil granules can be mobilized without the prior degranulation of all peroxidase-negative/specific granules. Accurate measurements of the release of distinct granules would help clarify this issue better. Attempts have been made but the technique used has not the resolution required and the results obtained have (according to the authors) to be considered with caution [15].

Taken together, our data reveal no granule subsets differing in calcium affinities, making the concept of hierarchical exocytosis based on this property not applicable. The results presented here differ in certain respects from earlier published data regarding the basic regulatory mechanisms in neutrophil granule secretion [12, 13, 15, 16]. Several factors can underlie these discrepancies. For example, the studies were performed in different cell types, different methods were used to elevate  $[Ca^{2+}]_i$  and/or induce secretion, and the duration of the measurements was also different. More importantly, the basic finding that not all specific granules must be released before secretion of azurophil granules may commence also finds support in previous investigations

[13, 27]. The precise mechanism(s) underlying the different kinetics of the responses is not clear at this time, although both surface-to-mass ratio [24] or differential expression of SNARE proteins [7–9] are attractive alternatives. Furthermore, also the membrane tension of a vesicle, being higher in smaller vesicles and lower in larger vesicles, may play a part in the process. Studies of fusion events occurring in artificial membrane systems indicate a similar behavior in that the smaller membrane vesicles are undergoing fusion faster, as compared to larger vesicles [28]. As such, the mere size of the vesicle may serve as a control element of the fusion event. In all cases, more calcium makes more granules eligible for release faster, resulting in a faster kinetics at higher  $[Ca^{2+}]_i$ .

In living cells, the cytoskeleton is essential for the control of activity and function [17, 29]. Therefore, data obtained and conclusions drawn from experimental systems with disrupted cytoskeleton (eg, systems using cytochalasin B to increase secretion) are, in our view, difficult to relate to the actual in vivo situation. In addition, we found it exceedingly hard to achieve good seals and whole-cell recordings in the presence of cytochalasin B. We were thus not able to investigate the direct effects of cytochalasin B on the secretory process under study.

In summary, we were unable to detect biphasic capacitance increases as has previously been described and suggested to reflect distinct high- as well as low-affinity granule

populations in human neutrophils. These findings extend and modulate previous knowledge regarding neutrophil granule mobilization. Recent studies on both eosinophils and neutrophils also reveal several distinct types of granule fusion events including not only fusion of single granules with the plasma membrane but also intracellular granule-granule fusion and fusion of large compounds of prefused granules with the plasma membrane [30]. It is obvious that such a pre-fusion process may involve granules of different types [31] and this may have implication for the data suggesting that not all specific granules have to be mobilized before release of azurophil granules will commence. The coexistence of distinct modes of exocytosis in which granule content from different granule subsets interact may be a mechanism for efficient targeting of release during exocytosis and further tune neutrophil function in the inflammatory response.

## ACKNOWLEDGMENTS

We thank Marie Samuelsson for skilful technical assistance. The staff at the Blood Donor Centre at SU/Sahlgrenska is acknowledged for their assistance. This study was supported by Grants from the Swedish Medical Research Council and King Gustaf the V 80-year Foundation.

## REFERENCES

- [1] Burg ND, Pillinger MH. The neutrophil: function and regulation in innate and humoral immunity. *Clinical Immunology*. 2001;99(1):7–17.
- [2] Berton G. Degranulation. In: Gallin JI, Snyderman R, eds. *Inflammation: Basic Principles and Clinical Correlates*. Philadelphia, Pa: Lippincott Williams & Wilkins; 1999:703–719.
- [3] Borregaard N, Cowland JB. Granules of the human neutrophilic polymorphonuclear leukocyte. *Blood*. 1997;89(10):3503–3521.
- [4] Sengelov H, Follin P, Kjeldsen L, Lollike K, Dahlgren C, Borregaard N. Mobilization of granules and secretory vesicles during in vivo exudation of human neutrophils. *Journal of Immunology*. 1995;154(8):4157–4165.
- [5] Sollner T, Whiteheart SW, Brunner M, et al. SNAP receptors implicated in vesicle targeting and fusion. *Nature*. 1993;362(6418):318–324.
- [6] Chen YA, Scheller RH. SNARE-mediated membrane fusion. *Nature Reviews Molecular Cell Biology*. 2001;2(2):98–106.
- [7] Brumell JH, Volchuk A, Sengelov H, et al. Subcellular distribution of docking/fusion proteins in neutrophils, secretory cells with multiple exocytic compartments. *Journal of Immunology*. 1995;155(12):5750–5759.
- [8] Martin-Martin B, Nabokina SM, Blasi J, Lazo PA, Mollinedo F. Involvement of SNAP-23 and syntaxin 6 in human neutrophil exocytosis. *Blood*. 2000;96(7):2574–2583.
- [9] Mollinedo F, Martin-Martin B, Calafat J, Nabokina SM, Lazo PA. Role of vesicle-associated membrane protein-2, through Q-soluble N-ethylmaleimide-sensitive factor attachment protein receptor/R-soluble N-ethylmaleimide-sensitive factor attachment protein receptor interaction, in the exocytosis of specific and tertiary granules of human neutrophils. *Journal of Immunology*. 2003;170(2):1034–1042.
- [10] Burgoyne RD, Morgan A.  $Ca^{2+}$  and secretory-vesicle dynamics. *Trends in Neurosciences*. 1995;18(4):191–196.
- [11] Andrews NW. Regulated secretion of conventional lysosomes. *Trends in Cell Biology*. 2000;10(8):316–321.
- [12] Lew PD, Monod A, Krause K-H, Waldvogel FA, Biden TJ, Schlegel W. The role of cytosolic free calcium in the generation of inositol 1,4,5-trisphosphate and inositol 1,3,4-trisphosphate in HL-60 cells. Differential effects of chemotactic peptide receptor stimulation at distinct  $Ca^{2+}$  levels. *Journal of Biological Chemistry*. 1986;261(28):13121–13127.
- [13] Barrowman MM, Cockcroft S, Gomperts BD. Differential control of azurophilic and specific granule exocytosis in Sendai-virus-permeabilized rabbit neutrophils. *Journal of Physiology*. 1987;383:115–124.
- [14] Borregaard N, Lollike K, Kjeldsen L, et al. Human neutrophil granules and secretory vesicles. *European Journal of Haematology*. 1993;51(4):187–198.
- [15] Nusse O, Serrander L, Lew DP, Krause K-H.  $Ca^{2+}$ -induced exocytosis in individual human neutrophils: high- and low-affinity granule populations and submaximal responses. *EMBO Journal*. 1998;17(5):1279–1288.
- [16] Nusse O, Lindau M. The calcium signal in human neutrophils and its relation to exocytosis investigated by patch-clamp capacitance and Fura-2 measurements. *Cell Calcium*. 1993;14(4):255–269.
- [17] Rothe G, Klouche M. Phagocyte function. *Methods in Cell Biology*. 2004;75:679–708.
- [18] Böyum A. Isolation of mononuclear cells and granulocytes from human blood. Isolation of mononuclear cells by one centrifugation, and of granulocytes by combining centrifugation and sedimentation at 1 g. *Scandinavian Journal of Clinical and Laboratory Investigation, Supplement*. 1968;97:77–89.
- [19] Hamill OP, Marty A, Neher E, Sakmann B, Sigworth FJ. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Archiv*. 1981;391(2):85–100.
- [20] Lindau M, Neher E. Patch-clamp techniques for time-resolved capacitance measurements in single cells. *Pflügers Archiv*. 1988;411(2):137–146.
- [21] Neher E, Marty A. Discrete changes of cell membrane capacitance observed under conditions of enhanced secretion in bovine adrenal chromaffin cells. *Proceedings of the National Academy of Sciences of the United States of America*. 1982;79(21):6712–6716.
- [22] Joshi C, Fernandez JM. Capacitance measurements. An analysis of the phase detector technique used to study exocytosis and endocytosis. *Biophysical Journal*. 1988;53(6):885–892.
- [23] Bers DM, Patton CW, Nuccitelli R. A practical guide to the preparation of  $Ca^{2+}$  buffers. *Methods in Cell Biology*. 1994;40:3–29.
- [24] Borregaard N. Development of neutrophil granule diversity. *Annals of the New York Academy of Sciences*. 1997;832:62–68.
- [25] Lollike K, Lindau M. Membrane capacitance techniques to monitor granule exocytosis in neutrophils. *Journal of Immunological Methods*. 1999;232(1-2):111–120.
- [26] Nusse O, Lindau M. The dynamics of exocytosis in human neutrophils. *Journal of Cell Biology*. 1988;107(6 I):2117–2123.
- [27] Sengelov H, Kjeldsen L, Borregaard N. Control of exocytosis in early neutrophil activation. *Journal of Immunology*. 1993;150(4):1535–1543.
- [28] Cans A-S, Wittenberg N, Karlsson R, et al. Artificial cells: unique insights into exocytosis using liposomes and lipid nanotubes. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;100(2):400–404.

- 
- [29] Granfeldt D, Dahlgren C. An intact cytoskeleton is required for prolonged respiratory burst activity during neutrophil phagocytosis. *Inflammation*. 2001;25(3):165–169.
- [30] Lollike K, Lindau M, Calafat J, Borregaard N. Compound exocytosis of granules in human neutrophils. *Journal of Leukocyte Biology*. 2002;71(6):973–980.
- [31] Karlsson A, Dahlgren C. Assembly and activation of the neutrophil NADPH oxidase in granule membranes. *Antioxidants and Redox Signaling*. 2002;4(1):49–60.



**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

