

# Trials of pharmacological interventions for Tourette Syndrome: A systematic review

Karen Waldon<sup>a</sup>, Jonathan Hill<sup>a</sup>, Cristiano Termine<sup>b</sup>, Umberto Balottin<sup>c</sup> and Andrea Eugenio Cavanna<sup>a,d,\*</sup>

<sup>a</sup>*The Michael Trimble Neuropsychiatry Research Group, Department of Neuropsychiatry, BSMHFT and University of Birmingham, Birmingham, UK*

<sup>b</sup>*Child Neuropsychiatry Unit, Department of Clinical and Experimental Medicine, University of Insubria, Varese, Italy*

<sup>c</sup>*Child Neuropsychiatry Unit, IRCCS 'C. Mondino Institute of Neurology' Foundation, University of Pavia, Pavia, Italy*

<sup>d</sup>*Department of Motor Neuroscience and Movement Disorders, Institute of Neurology and University College London, London, UK*

**Abstract.** *Introduction:* Gilles de la Tourette Syndrome (GTS) is a childhood-onset hyperkinetic movement disorder defined by the chronic presence of multiple motor tics and at least one vocal tic and often complicated by co-morbid behavioural problems. The pharmacological treatment of GTS focuses on the modulation of monoaminergic pathways within the cortico-striato-thalamo-cortical circuitry. This paper aims to evaluate the efficacy and safety profiles of pharmacological agents used in the treatment of tics in patients with GTS, in order to provide clinicians with an evidence-based rationale for the pharmacological treatment in GTS.

*Method:* In order to ascertain the best level of evidence, we conducted a systematic literature review to identify double-blind randomised controlled trials of medications in GTS populations.

*Results:* We identified a large number of pharmacological agents as potentially effective in improving tic symptoms. The alpha-2 agonist Clonidine is amongst the agents with the most favourable efficacy-versus-adverse events ratio, especially in patients with co-morbid attention-deficit hyperactivity disorder, although effect sizes vary evidence-based studies.

*Discussion:* Our results are in line with the findings of uncontrolled open-label studies. However, most trials have low statistical power due to the small sample sizes, and newer agents, such as Aripiprazole, have not been formally tested in double-blind randomised controlled trials. Further research should focus on better outcome measures, including Quality of Life instruments.

**Keywords:** Drug therapy, movement disorder, pharmacological treatment, randomised controlled trial, tics, Tourette Syndrome

## 1. Introduction

Gilles de la Tourette Syndrome (GTS) is a childhood-onset neurological disorder [1] characterised by repetitive, stereotypical involuntary movements and vocalisations known as tics [2], which vary in severity over weeks to months [3]. GTS is often associated with be-

havioural problems, such as attention deficit hyperactivity disorder (ADHD) and obsessive-compulsive disorder (OCD) [4,5]. It is estimated that GTS affects 1–8 per 1000 children [6] worldwide [7], in different degrees of severity [7], however due to misdiagnosis in reality prevalence may be even higher.

Current treatment options for GTS include behavioural therapy [8], medication [9] and neurosurgery [10]. Examples of psychobehavioural interventions are exposure and response prevention [11], massed practice and habit reversal training [11], as well as relaxation therapy [11], biofeedback [11] and acupuncture [12].

---

\*Correspondence: Dr. Andrea Eugenio Cavanna MD, Department of Neuropsychiatry, The Barberry National Centre for Mental Health, B15 2FG, Birmingham, UK. Tel.: +44 121 3012317; Fax: +44 121 3012291; E-mail: a.cavanna@ion.ucl.ac.uk.




Fig. 1. Flow chart summarising the study selection process.

Medications include typical antipsychotics (neuroleptics) [13,14], atypical antipsychotics [15–17], alpha-adrenergic agonists [18–20], along with other pharmacological classes [21–27].

Historically, neuroleptics such as haloperidol and pimozide have been the mainstay of treatment [4,28], however due to their metabolic and extrapyramidal side effects [28–30] focus has more recently shifted to other agents, such as atypical antipsychotics [15,16,31,32], which have variable D2 receptor blocking properties and 5-HT receptor blocking properties [33]. These however can also cause metabolic side effects, such as weight gain [15,29,31], which increases risk of cardiovascular disease and type 2 diabetes [27,34].

As a relatively common condition, GTS can cause disruption at school and embarrassment for the child [35] subsequently impacting on their education [20] in addition to difficulties with social interactions [20]. Improving pharmacological control over tics could alleviate these problems [9] but may cause adverse effects of their own [33], outweighing the benefits.

In this paper, we set out to systematically review the existing evidence on medications used to treat GTS. Our aim was to evaluate these agents' efficacy (reduction in tic frequency and severity as well as overall impairment), weighed up against risk of adverse effects, in order to inform clinical practice about the evidence-based rationale for the choice of the most appropriate treatment and ultimately provide better care for this patient population.

## 2. Methods

We conducted a systematic review of the available literature, with focus on randomised, controlled, double-

blind studies comparing pharmacological therapies to placebo for the treatment of GTS. Both parallel group and cross-over study designs were included.

Our electronic searches were conducted on the databases Medline and PsycINFO. The first search cross-referenced “Tourette” and its derivations with drug names, as MeSH headings and as text words in the title and abstract fields. Both search terms were required to be major descriptors. The same cross-referenced search was carried out with “tic” in place of “Tourette”. Type of article was restricted to original trials on human subjects, published in peer-reviewed journals. Only articles published in the English language since 1960 were included. Since GTS is a disorder which first presents in childhood, we included both children and adult studies.

Relevant journals were manually searched for additional articles using indexes or online search engines. These journals included *Neurology*, *American Journal of Psychiatry*, *Journal of Neuropsychiatry and Clinical Neurosciences*, *Movement Disorders*, *Annals of Neurology*, *Brain*, *Journal of Neurology*, *Neurosurgery and Psychiatry* and *Journal of Psychosomatic Research*. Finally, the reference lists of relevant articles were manually searched for related studies.

The steps of our search methodology are summarised in Fig. 1.

## 3. Results

Thirty-three studies (17 parallel trials and 16 crossover trials) involving a total of 1,385 patients fulfilled our inclusion criteria and were included in the

Table 1

Double-blind randomised controlled studies investigating the efficacy of antipsychotic medications for tic management in Gilles de la Tourette syndrome

Medication	Author	Year	Design	N	Age range (years)	Treatment duration (weeks)	Effect on tics	Advantages	Disadvantages	
<i>Typical antipsychotics</i>										
Haloperidol	Silver et al.	2001	Parallel	70		6.5	+		SE: Some cases of nausea and vomiting	
	Ross and Moldofsky	1978	Parallel	9		80	+			
	Sallee et al.	1996	Crossover	26	7–13	6	+		SE: 3-fold higher frequency of serious SE and greater ESE than Pimozide SE common SE: Sedation and Weight Gain	
	Sallee et al.	1997	Crossover	22	7–16	6	–			
	Pimozide	Shapiro et al.	1989	Parallel	57			+		
		Bruggeman et al.	2001	Parallel	41	11–65	7	+		ESE rare
	Pimozide	Gilbert et al.	2004	Crossover	13	7–17	4	+	No associated ECG changes Less weight gain than Risperidone Less lethargy than Haloperidol	SE: Prolonged QTc interval
		Ross and Moldofsky	1978	Parallel	9		80	+		
		Sallee et al.	1996	Crossover	26	7–13	6	+		
		Sallee et al.	1997	Crossover	22	7–16	6	+	Less SE than Haloperidol	
Shapiro et al.		1989	Parallel	57			+			
Shapiro et al.		1984	Crossover	20		6	+			
Atypical antipsychotics	Risperidone	Onofrj et al.	2000	Crossover	4	19–40	52	+	SE: Sedation and minor motor SE	
		Bruggeman et al.	2001	Parallel	41	11–65	7	+	ESE rare	
	Risperidone	Gilbert et al.	2004	Crossover	13	7–17	4	+	More effective than Pimozide	SE: Sedation and Weight Gain SE: Greater weight gain than Pimozide
		Dion et al.	2002	Parallel	48		8	+	No increase in OCS	SE: Fatigue, somnolence, hypokinesia, tremor
		Scahill et al.	2003	Parallel	34	6–62	8	+	No ESE	SE: Acute social phobia, weight gain
		Gaffney et al.	2002	Parallel	21	7–17	8	+	No ESE	SE: Sedation
	Olanzapine	Onofrj et al.	2000	Crossover	4	19–40	52	+	More effective than Pimozide	SE: Minor sedation

Abbreviations: N = number of subjects in the study; SE = Side Effects; ESE; Extrapyramidal Side Effects; OCS = Obsessive Compulsive symptoms; TSSG = Tourette's Syndrome Study Group.

present review. Table 1 summarises the double-blind randomised controlled studies investigating the efficacy of typical [14,30] and atypical antipsychotic medications [15–17,31,32] for tic management in GTS, whilst Table 2 focuses on the non-antipsychotic medications: alpha2 adrenergic agonists [18–20,36], anticonvulsants [23], dopamine agonists [24,36,37], GABA-

modulating agents [21], substituted benzamides [5,38], anticholinergic agents [26], cannabinoids [27], antiemetics [25,39], 5HT3 antagonists [25], Selective Serotonin Reuptake Inhibitors (SSRIs) [5,40], Selective Norepinephrine Reuptake Inhibitors (SNRIs) [41], tricyclic antidepressants [42], MAO inhibitors and antian- drogens [43,44].

Table 2

Double-blind randomised controlled studies investigating the efficacy of non-antipsychotic medications for tic management in Gilles de la Tourette syndrome

Medication	Author	Year	Design	N	Age range (years)	Treatment duration (weeks)	Effect on tics	Advantages	Disadvantages
<i>Alpha adrenergic agonists</i>									
Clonidine	Goetz et al.	1987	Crossover	30			–		
	Singer et al.	1995	Crossover	34	7–13	6	–	Helped with ADHD symptoms	SE: Sedation
	Hedderick et al.	2009	Crossover	10	8–27	6	+		SE: Sedation
	Gaffney et al.	2002	Parallel	21	7–17	8	+	No ESE	SE: Sedation
	Du et al.	2008	Parallel	437	6–18	4	+	No SE	
	TSSG	2002	Parallel	103	7–14	16	+	Helped with ADHD symptoms	SE: Sedation
Guanfacine	Scahill et al.	2001	Parallel	34	6–14	8	+	Helped with ADHD symptoms	SE: Sedation
<i>Anticonvulsants</i>									
Topiramate	Jankovic	2009	Parallel	20	7–65		+	No observed adverse SE.	
Levetiracetam	Hedderick et al.	2009	Crossover	10	8–27	6	–		SE: Irritability
	Smith-Hicks et al.	2007	Crossover	22	9–14	4	–		
<i>Dopamine agonists</i>									
Pergolide	Gilbert et al.	2000	Crossover	24	7–17	6	+	Well tolerated and no serious SE.	
	Gilbert et al.	2003	Parallel	57	7–17		+	Well tolerated, helps with ADHD symptoms	
Talipexole	Goetz et al.	1994	Parallel	13			–		SE: Sedation, dizziness
<i>GABA modulating agents</i>									
Baclofen	Singer et al.	2001	Crossover	9		4	+	No major side effects	YGTSS scores showed reduction in impairment scores rather than decrease in tics
<i>Substituted benzamides</i>									
Sulpiride	George et al.	1993	Crossover	11			+		
Tiapride	Eggers et al.	1988	Crossover	17			+	Minimal SE: hyperprolactinaemia restricted to duration of therapy	
<i>Acetylcholinergic agents</i>									
Nicotine	Silver et al.	2001	Parallel	70		2.5	+	Combination with haloperidol superior to haloperidol only	SE: Nausea and Vomiting common
Mecamylamine	Silver, Sheehan et al.	2001	Parallel	38	8–17	8	–	Reduction in sudden mood changes and depression	
<i>Cannabinoids</i>									
Delta-9-tetrahydrocannabinol	Muller-Vahl et al.	2003	Parallel	17		6	+	No serious adverse SE	
<i>Anti-emetics</i>									
Metoclopramide	Nicholson et al.	2005	Parallel	27	9–14	8	+	Well tolerated, no significant cardiac changes. Small increase in serum Prolactin	Further trials needed in younger patients

Table 2, continued

Medication	Author	Year	Design	N	Age range (years)	Treatment duration (weeks)	Effect on tics	Advantages	Disadvantages
<i>5HT3 Antagonists</i>									
Ondansetron	Toren et al.	2005	Parallel	30	12–46	3	–		SE: Abdominal pain
<i>SSRIs</i>									
Fluoxetine	Seahill et al.	1997	Crossover	14	8–33	8	–	Reduction in OCS	SE: Transient behavioural activation
<i>SNRIs</i>									
Atomoxetine	Allen et al.	2005	Parallel	148	7–17	18	+	Useful for ADHD symptoms	SE: Nausea, decreased appetite
<i>Tricyclic anti-depressants</i>									
Desipramine	Singer et al.	1995	Crossover	34	7–13	6	–	Helped with ADHD symptoms	
Fluvoxamine	George et al.	1993	Crossover	11			–		
<i>MAO inhibitors</i>									
Deprenyl	Feigin et al.	1996	Crossover	15	7–16	8	+	No SE	
<i>Antiandrogens</i>									
Flutamide	Peterson et al.	1998	Crossover	13		3	+/-	Modest improvement in OCS	Effect is short-lived
							(reduction in motor but not phonic tics)		

–+ = Positive effect on tic severity; – = Evidence of poor treatment response

Abbreviations: N = number of subjects in the study; SE = Side Effects; ESE; Extrapyrimal Side Effects; OCS = Obsessive Compulsive symptoms; TSSG = Tourette's Syndrome Study Group.

With regards to outcome measures, the majority of trials used Clinical Global Impression (CGI) scale [19,45] and the Yale Global Tic Severity Scale (YGTSS) [45] as the main outcome measure [21,23,24,27,46–48]. The YGTSS is the total of 2 subscale scores; total tic score (TTS) which rates tic severity from 0–5 and tic impairment score (TIS) which represents how GTS impacts for example family life and self-esteem [23]. The CGI scale rates how much GTS affects a patient's life [23]. Other measures include the Tourette's Disorder Scale-Clinician Rated (TODS-CR) [26] and video tapings used to assess tic frequency by clinicians [36,38,49].

#### 4. Discussion

In this paper, we reviewed the existing evidence from randomised double-blind controlled trials investigating the pharmacological treatment of tics in GTS. We critically evaluated efficacy in terms of reduction in tic severity and presence of adverse effects, with the aim to inform clinical practice regarding the choice of pharmacotherapy in GTS.

Antipsychotic medications have long been used for the treatment of tics in GTS [30]. Consistent with early findings, typical antipsychotics Haloperidol and

Pimozide demonstrated significant efficacy (reduction in tic severity) [13,14,16,28,30,31,50], however they were also characterised by common side effects, such as sedation [17], weight gain, depression [16], extrapyramidal symptoms [17,30] (especially Haloperidol [28,30]) and risk of QTc prolongation [14,51] (especially Pimozide [14]). According to the available evidence, these agents should not be recommended as first line medications, due to the high rate of adverse effects [28,29]. Among the newer antipsychotics, the use of Risperidone is arguably supported by the best evidence for efficacy [15,16,28,31,32,52,53]. However, weight gain [15,31] and sedation [53] have been reported as common side effects [32], followed by hyperprolactinaemia and risk of extrapyramidal symptoms [16,32]. The available evidence suggests that Risperidone may not be suitable for overweight patients [28,31], whilst it can prove beneficial for patients with comorbid OCD [16,53] as augmentation therapy to SSRIs for treatment-refractory obsessive-compulsive symptoms. The same is true for substituted benzamides sulpiride [5] and tiapride [5,54–58], although reports of tardive dyskinesia following use of sulpiride use warrant caution [59]. Little evidence was available for new generation antipsychotic medications, such as D2 partial agonist Aripiprazole, which has been recently shown to have positive efficacy and tolerability profiles

in a series of case reports and open label studies [51, 60,61].

The centrally acting alpha 2-adrenergic agonist Clonidine [18] has been shown to be a potentially effective and safe option for the first-line treatment of tics in GTS, although effect sizes showed a degree of variability across different studies [3,20,47,53,54,62]. Several studies also reported a significant improvement for patients with comorbid ADHD in impulsive-hyperactive symptoms [19] as well as tic symptoms with both Clonidine [42] and Guanfacine [19]. Among other agents, the newer antiepileptic drugs deserve attention in consideration of the number of trials carried out in GTS populations in recent years. In particular, Topiramate was found to be an effective medication for tic reduction [4], especially in patients suffering from obesity or migraine. However, previous research also reported relatively frequent side effects associated with its use, including concentration difficulties, confusion, ataxia and diplopia [63–65]. The current evidence for Lev- etiracetam is somewhat conflicting, with a number of studies showing efficacy [66–69], and others showing Lev- etiracetam being less effective than Clonidine and with no significant improvement in tic severity [23,47]. Moreover, adverse reactions to this medication include irritability, hyperkinesias and insomnia [47].

The reviewed literature presents a number of limitations, which might have a significant impact on the quality of the results we obtained. One main limitation of the trials reviewed in this paper is the high rate of patients lost at follow-up [26], which can be as high as 55% [5], thus reducing the reliability of the results. However ad hoc analysis showed few differences between whole group and group minus drop-outs in endpoint [3,5]. The majority of patients were recruited from specialist clinics; therefore the sample may not be representative of the general population because of the possibility of referral bias. Some of the studies identified were rather small in terms of numbers of participants [17,18,36,40,47]. Small sample sizes mean a relatively small effect in a few patients can result in large treatment effects, which can reduce the reliability of the results. Some studies excluded patients with co-morbid OCD and ADHD, thus reducing the sample size and making the sample less representative. On the other hand, the presence of behavioural co-morbidities can influence the study findings, if not adequately accounted for. Some studies did not provide any information on whether participants had ceased medications (or had a wash-out period) before starting the study, thus leading to potential drug-drug interactions

as additional confounding factors [21]. Likewise, the method of randomisation was not reported in all trials. Despite randomisation, there were baseline differences between groups, which in some studies were accounted for using analysis of covariance [25]. Although the results of some studies showed that these differences did not affect the results [37,46], more severe tics at baseline can lead to numerically greater improvements than milder tics [21]. Studies also varied in their approach to determine treatment efficacy, sometimes making it difficult to compare and standardise their results. For example, certain outcome measures such as clinician-assessed video recordings [18,36,49] increase the risk of measurement error and bias. However, most trials were assessed by multiple clinicians and few discrepancies were found between their ratings, indicating a good degree of reliability. Although a number of trials reported a statistical significance in ratings of tic severity, clinical significance was not always apparent. Finally, a mixture of crossover and parallel studies were reviewed, which can have caused some inconsistency in the interpretation of the findings.

Future studies investigating the pharmacological treatment of GTS should be conducted with focus of newly developed, promising agents belonging to pharmaceutical classes with proven efficacy [39], such as the newer antipsychotic Aripiprazole [51,60,61]. With regards to known treatments, further randomised placebo-controlled double-blinded studies are needed with greater sample sizes to increase statistical power, and lower loss to follow up. These studies should ideally use multidimensional outcome measures, such as Quality of Life (QoL) ratings [70], alongside tic severity ratings. Recent research has shown that there is not always a direct correlation between tic severity and QoL, implicating the usefulness of QoL measures in treatment trials. Other primary outcome measures should include ratings of symptoms of co-morbid ADHD and OCD, to address the holistic treatment of patients with GTS. Intention-to-treat analysis should be performed to monitor adverse effects. Finally, samples should also be recruited from different community and clinic populations (e.g. by advertisements in mainstream schools, as well as special education schools and specialist clinics).

## 5. Conclusion

The aim of this paper was to inform clinicians about the appropriate pharmacological management of pa-

tients with GTS based on the available evidence. We conducted a systematic literature review focusing on high-quality randomised placebo-controlled trials by using strict search criteria. A number of pharmacological agents were found to be potentially effective in improving tic severity. However some medications (e.g. typical antipsychotics) were characterised by a less favourable profile due to the presence of significant side-effects [46], whilst other agents (e.g. atypical antipsychotics; alpha-2 adrenergic agonists, especially Clonidine [42,71]) showed added benefits of treating co-morbid behavioural symptoms [19,40,41, 43,48]. Caution should be used to avoid the risk of misinterpreting the findings of these studies, as for instance effect sizes show considerable variation in the case of Clonidine. Moreover, the lack of high-level evidence trials makes it difficult to derive conclusions about other promising agents, such as the D2 partial agonist Aripiprazole [51,60,61]. In consideration of the major limitations of the currently available literature (most studies have been conducted on small samples and their results have not been robustly replicated), further research is needed to establish an evidence-based rationale for the choice of first- and second-line pharmacological options for the treatment of tics in patients with GTS.

## Acknowledgements

The authors are grateful to Tourettes Action UK for their continuing support.

## References

- [1] World Health Organisation. *The International Classification of Diseases*, 10th ed., Geneva, 1992.
- [2] A.E. Cavanna, S. Servo, F. Monaco and M.M. Robertson, The behavioural spectrum of Gilles de la Tourette Syndrome, *J Neuropsychiatry Clin Neurosci* **21**(1) (2009), 13–23.
- [3] J.F. Leckman, M.T. Hardin, M.A. Riddle, J. Stevenson, S.I. Ort and D.J. Cohen, Clonidine treatment of Gilles de la Tourette's syndrome, *Arch Gen Psychiatry* **48**(4) (1991), 324–328.
- [4] J. Jankovic, J. Jimenez-Shahed and L.W. Brown, A randomised, double-blind placebo-controlled study of topiramate in the treatment of Tourette Syndrome, *J Neurol Neurosurg Psychiatry* **81** (2010), 70–73.
- [5] M.S. George, M.R. Trimble and M.M. Robertson, Fluvoxamine and Sulpiride in Comorbid Obsessive-Compulsive Disorder and Gilles de la Tourette Syndrome, *Hum Psychopharmacol* **8**(5) (1993), 327–334.
- [6] B. Kadesjo and C. Gillberg, Tourette's disorder: epidemiology and comorbidity in primary school children, *J Am Acad Child Adolesc Psychiatry* **39**(5) (2000), 548–555.
- [7] M.M. Robertson, V. Eapen and A.E. Cavanna, The international prevalence, epidemiology, and clinical phenomenology of Tourette Syndrome: a cross-cultural perspective, *J Psychosom Res* **67**(6) (2009), 475–483.
- [8] M.M. Robertson, Gilles de la Tourette syndrome: the complexities of phenotype and treatment, *Br J Hosp Med (Lond)* **72**(2) (2011), 100–107.
- [9] C.M. Eddy, H.E. Rickards and A.E. Cavanna, Treatment strategies for tics in Tourette Syndrome, *Ther Adv Neurol Disord* **4**(1) (2011), 25–25.
- [10] M. Porta, M. Sassi, F. Ali, A.E. Cavanna and D. Servello, Neurosurgical treatment for Gilles de la Tourette syndrome: the Italian perspective, *J Psychosom Res* **67**(6) (2009), 585–590.
- [11] C. Verdellen, J. van de Griendt, A. Hartmann and T. Murphy, European clinical guidelines for Tourette syndrome and other tic disorders. Part III: behavioural and psychosocial interventions, *Eur Child Adolesc Psychiatry* **20**(4) (2011), 197–207.
- [12] L. Wu, H. Li and L. Kang, 156 cases of Gilles de la Tourette's syndrome treated by acupuncture, *J Tradit Chin Med* **16**(3) (1996), 211–213.
- [13] M.S. Ross and H. Modolfsky, A comparison of pimozide and haloperidol in the treatment of Gilles de la Tourette's syndrome, *Am J Psychiatry* **135**(5) (1978), 585–587.
- [14] E. Shapiro, A.K. Shapiro, G. Fulop, M. Hubbard, J. Mandeli, J. Nordlie and R.A. Phillips, Controlled study of haloperidol, pimozide and placebo for the treatment of Gilles de la Tourette's syndrome, *Arch Gen Psychiatry* **46**(8) (1989), 722–730.
- [15] L. Scahill, J.F. Leckman, R.T. Schultz, L. Katsovich and B.S. Peterson, A Placebo-controlled trial of risperidone in Tourette Syndrome, *Neurology* **60**(7) (2003), 1130–1135.
- [16] R. Bruggeman, C. van der Linden, J.K. Buitelar, G.S. Gericke, S.M. Hawkrige and J.A. Temlett, Risperidone versus pimozide in Tourette's disorder: a comparative double-blind parallel-group study, *J Clin Psychiatry* **62**(1) (2001), 50–56.
- [17] M. Onofrij, C. Paci, G. D'Andreammateo and L. Toma, Olanzapine in severe Gilles de la Tourette syndrome: a 52 week double blind crossover study vs. low-dose pimozide, *J Neurol* **247**(6) (2000), 443–446.
- [18] C.G. Goetz, C.M. Tanner, R.S. Wilson, V.S. Carrol, P.G. Como and K.M. Shannon, Clonidine and Gilles de la Tourette syndrome: double-blind study using objective rating methods, *Ann Neurol* **21**(3) (1987), 307–310.
- [19] L. Scahill, P.B. Chappell, Y.S. Kim, R.T. Schultz, L. Katsovich, E. Shepherd, A.F. Arnsten, D.J. Cohen and J.F. Leckman, A placebo-controlled study of guanfacine in the treatment of children with tic disorders and attention deficit hyperactivity disorder, *Am J Psychiatry* **158**(7) (2001), 1067–1074.
- [20] Y. Du, H. Li, A. Vance, Y. Zhong, F. Jiao, H. Wang, M. Wang, L. Su, D. Yu, S. Ma and J. Wu, Randomized double-blind multicentre placebo-controlled clinical trial of the clonidine adhesive patch for the treatment of tic disorders, *Aust N Z J Psychiatry* **42**(9) (2008), 807–813.
- [21] H.S. Singer, J. Wendlandt, M. Krieger and J. Giulliano, Baclofen Treatment in Tourette syndrome: A double-blind, placebo-controlled, crossover trial, *Neurology* **56**(5) (2001), 599–604.
- [22] M.M. Robertson and J.S. Stern, Gilles de la Tourette Syndrome: symptomatic treatment based on evidence, *Eur Child Adolesc Psychiatry* **9**(1) (2000), 60–75.
- [23] C.L. Smith-Hicks, D.D. Bridges, N.P. Paynter and H.S. Singer, A Double Blind Randomized Placebo Control Trial of Levitiracetam in Tourette Syndrome, *Mov Disord* **22**(12) (2007), 1764–1770.

- [24] D.L. Gilbert, G. Sethuraman, L. Sine, S. Peters and F.R. Sallee, Tourette's syndrome improvement with pergolide in a randomized, double-blind, crossover trial, *Neurology* **54**(6) (2000), 1310–1315.
- [25] P. Toren, A. Weizman, S. Ratner, D. Cohen and N. Laor, Ondansetron treatment in Tourette's disorder: A 3-week, randomized, double-blind, placebo-controlled study, *J Clin Psychiatry* **66**(4) (2005), 499–503.
- [26] A.A. Silver, R.D. Shytle, K.H. Sheehan, D.V. Sheehan, A. Ramos and P.R. Sanberg, Multicentre, double-blind, placebo-controlled study of mecamylamine monotherapy for Tourette's disorder, *J Am Acad Child Adolesc Psychiatry* **40**(9) (2001), 1103–1110.
- [27] K.R. Muller-Vahl, U. Schneider, H. Prevedel, K. Theloe, H. Kolbe, T. Daldrup and H.M. Emrich, Delta 9-tetrahydrocannabinol (THC) is effective in the treatment of tics in Tourette Syndrome: a 6-week randomized trial, *J Clin Psychiatry* **64**(4) (2003), 459–465.
- [28] F.R. Sallee, L. Nesbitt, C. Jackson, L. Sine and G. Sethuraman, Relative Efficacy of haloperidol and pimozide in children and adolescents with Tourette's disorder, *Am J Psychiatry* **154**(8) (1997), 1057–1062.
- [29] T. Pringsheim and C. Marras, Pimozide for tics in Tourette's Syndrome, *Cochrane Database Syst Rev* **15**(2) (2009), CD006996.
- [30] F.R. Sallee, D. Dougherty, G. Sethuraman and N. Vrindavanam, Prolactin monitoring of haloperidol and pimozide treatment in children with Tourette's Syndrome, *Biol Psychiatry* **40**(10) (1996), 1044–1050.
- [31] D.L. Gilbert, J.R. Batterson, G. Sethuraman and F.R. Sallee, Tic reduction with risperidone versus pimozide in a randomized, double-blind, crossover trial, *J Am Acad Child Adolesc Psychiatry* **43**(2) (2004), 206–214.
- [32] Y. Dion, L. Annable and P. Sandor, G. C. Risperidone in the treatment of Tourette Syndrome: A double-blind, placebo-controlled trial, *J Clin Psychopharmacol* **22**(1) (2002), 31–39.
- [33] A.O. Alao, K. Malhotra and M.J. Dewan, Comparing the side effect profile of the atypical antipsychotics, *West Afr J Med* **21**(4) (2002), 313–315.
- [34] National Institute for Health and Clinical Excellence. Clinical guideline CG43 Obesity, 2006.
- [35] C.M. Eddy, R. Rizzo, M. Gulisano, A. Agodi, M. Barchitta, P. Cali, M.M. Robertson, A.E. Cavanna, Quality of life in young people with Tourette Syndrome: a controlled study, *J Neurol* **258**(2) (2011), 291–301.
- [36] C.G. Goetz, G.T. Stebbins and J.A. Thelen, Talipexole and adult Gilles de la Tourette's Syndrome: Double-Blind, Placebo-Controlled Clinical Trial, *Mov Disord* **9**(3) (1994), 315–317.
- [37] D.L. Gilbert, L. Dure, G. Sethuraman, D. Raab, J. L. and F.R. Sallee, Tic reduction with pergolide in a randomized controlled trial in children, *Neurology* **60**(4) (2003), 606–611.
- [38] C. Eggers, A. Rothenberger and U. Berghaus, Clinical and neurobiological findings in children suffering from tic disease following treatment with tiapride, *Eur Arch Psychiatr Neurol Sci* **237**(4) (1988), 223–229.
- [39] R. Nicholson, B. Craven-Thuss, J. Smith, B.D. McKinlay, F.X. Castellanos, A randomised, double-blind placebo-controlled trial of Metoclopramide for the treatment of Tourette's disorder, *J Am Acad Child Adolesc Psychiatry* **44**(7) (2005), 640–646.
- [40] L. Scahill, M.A. Riddle, R.A. King, M.T. Hardin, A. Rasmussen, R.W. Makuch and J.F. Leckman, Fluoxetine has no marked effect on tic symptoms in patients with Tourette's syndrome: a double-blind placebo-controlled study, *J Child Adolesc Psychopharmacol* **7**(2) (1997), 75–85.
- [41] A.J. Allen, R.M. Kurlan, D.L. Gilbert, B.J. Coffey, S.L. Linder, D.W. Lewis, P.K. Winner, D.W. Dunn, L.S. Dure, F.R. Sallee, D.R. Milton, M.L. Mintz, R.K. Ricardi, G. Erenberg, L.L. Layton, P.D. Feldman, D.K. Kelsey and T.J. Spencer, Atomoxetine treatment in children and adolescents with ADHD and comorbid tic disorders, *Neurology* **65**(12) (2005), 1941–1949.
- [42] H.S. Singer, J. Brown, S. Quaskey, L.A. Rosenberg, E.D. Melits and M.B. Denckla, The treatment of attention-deficit hyperactivity disorder in Tourette's syndrome: a double-blind, placebo-controlled study with clonidine and desipramine, *Pediatrics* **95**(1) (1995), 74–81.
- [43] B.S. Peterson, H. Zhang, G. Anderson and J.F. Leckman, A Double-Blind, Placebo-Controlled, Crossover trial of an antiandrogen in the treatment of Tourette's Syndrome, *J Clin Psychopharmacol* **18**(4) (1998), 324–331.
- [44] B.S. Peterson, J.F. Leckman, L. Scahill and F. Naftolin, Steroid hormones and Tourette's syndrome: Early experience with antiandrogen therapy, *J Clin Psychopharmacol* **14**(2) (1994), 131–135.
- [45] J.T. Walkup, L.A. Rosenberg, J. Brown and H.S. Singer, The validity of instruments measuring tic severity in Tourette's syndrome, *J Am Acad Child Adolesc Psychiatry* **31**(3) (1992), 472–477.
- [46] A.A. Silver, R.D. Shytle, M.K. Philipp, B.J. Wilkinson, B. McConville and P.R. Sanberg, Transdermal Nicotine and Haloperidol in Tourette's Disorder: A double-blind placebo-controlled study, *J Clin Psychiatry* **62**(9) (2001), 707–714.
- [47] E.F. Hedderick, C.M. Morris and H.S. Singer, Double-blind, crossover study of clonidine and levetiracetam in Tourette syndrome, *Pediatr Neurol* **40**(6) (2009), 420–425.
- [48] A. Feigin, R. Kurlan, M.P. McDermott, J. Beach, T. Dimitropoulos, C.A. Brower, L. Chapieski, K. Trinidad, P. Como and J. Jankovic, A controlled trial of deprenyl in children with Tourette's syndrome and attention deficit hyperactivity disorder, *Neurology* **46**(4) (1996), 965–968.
- [49] A. Braun, M.M. Mouradian, E. Mohr, G. Fabbrini and T.N. Chase, Selective D-1 dopamine receptor agonist effects in hyperkinetic extrapyramidal disorders, *J Neurol Neurosurg Psychiatry* **52**(5) (1989), 631–635.
- [50] A.K. Shapiro and E. Shapiro, Controlled study of pimozide vs placebo in Tourette's syndrome, *J Am Acad Child Psychiatry* **23**(2) (1984), 161–173.
- [51] M. Gulisano, P.V. Cali, A.E. Cavanna, C. Eddy, H. Rickards and R. Rizzo, Cardiovascular safety of aripiprazole and pimozide in young patients with Tourette syndrome, *Neurol Sci* **32**(6) (2011), 1213–1217.
- [52] B. Kim, C. Lee, J. Hwang, M. Shin and S. Cho, Effectiveness and safety of risperidone for children and adolescents with chronic tic or Tourette disorders in Korea, *J Child Adolesc Psychopharmacol* **15**(2) (2005), 318–324.
- [53] G.R. Gaffney, P.J. Perry, B.C. Lund, K.A. Bever-Stille, S. Arndt and S. Kuperman, Risperidone versus clonidine in the treatment of children and adolescents with Tourette's syndrome, *J Am Acad Child Adolesc Psychiatry* **41**(3) (2002), 330–336.
- [54] I. Drtilkova, B. Balastikova and H. Lemanova, Tiapride, clonazepam and clonidine in children with tic and comorbid hyperkinetics disorder: Clinical EEG study, *Ceska Slov Psychiatr* **93**(3) (1997), 54–78.
- [55] I. Drtilkova, B. Balastikova, H. Lemanova and J. Zak, Clonazepam, clonidine and tiapride in children with tic disorder, *Homeost Health Dis* **37**(5) (1996), 216.



- [56] F.J. Jimenez-Jimenez and P.J. Garcia-Ruiz, Pharmacological Options for the treatment of Tourette's disorder, *Drugs* **61**(15) (2001), 2207–2220.
- [57] M.M. Robertson, V. Schnieden and A.J. Lees, Management of Gilles de la Tourette syndrome using Sulpiride, *Clin Neuropharmacol* **13**(3) (1990), 229–235.
- [58] C. Ho, H. Chen, N. Chiu, E. Shen and H. Lue, Short-term treatment of children and adolescents with Tourette syndrome or chronic tic disorder, *J Formos Med Assoc* **108**(10) (2009), 788–793.
- [59] V. Eapen, C.L. Katona, T.R. Barnes and M.M. Robertson, Sulpiride-induced tardive dyskinesia in a person with Gilles de la Tourette syndrome, *J Psychopharmacol* **7**(3) (1993), 290–292.
- [60] A.E. Cavanna, C. Selvini, C. Termine, C. Luoni, C.M. Eddy and H. Rickards, Tolerability profile of aripiprazole in patients with Tourette's syndrome, *J Psychopharmacol* 2011, Epub ahead of print.
- [61] K. Lewis, L. Rappa, D.A. Sherwood-Jachimowicz and M. Larosse-Pierre, Aripiprazole for the treatment of adolescent Tourette's syndrome: a case report, *J Pharm Pract* **23**(3) (2010), 239–244.
- [62] J.F. Leckman, J. Detlor, D.F. Harcherik, S. Ort, B.A. Shaywitz and D.J. Cohen, Short- and long-term treatment of Tourette's syndrome with clonidine: a clinical perspective, *Neurology* **35**(3) (1985), 343–351.
- [63] S. Kuo, J. Topiramate in treatment of Tourette syndrome, *Clin Neuropharmacol* **33**(1) (2010), 32–34.
- [64] T.Y. Nelson, M.T. Bost and P.S. Lesser, Topiramate in children and adolescents with Tourette syndrome: A case series, *Journal of Pediatric Neurology* **5**(1) (2007), 1304–2580.
- [65] F.S. Abuzzahab and L. Victoria, Control of Tourette's syndrome with topiramate, *Am J Psychiatry* **158**(6) (2001), 968.
- [66] Y. Awaad, A.M. Michon, S. Minarik and T. Rizk, Levitiracetam in Tourette syndrome: A randomized double blind, placebo controlled study, *Pediatr Neurol* **7**(3) (2009), 257–263.
- [67] Y. Awaad, A.M. Michon and S. Minarik, Long-term use of levitiracetam to treat tics in children and adolescents with Tourette syndrome, *Pediatr Neurol* **5**(3) (2007), 209–214.
- [68] Y. Awaad, A.M. Michon and S. Minarik, Use of levitiracetam to treat tics in children and adolescents with Tourette syndrome, *Mov Disord* **20**(6) (2005), 714–718.
- [69] A. Fernandez-Jaen, D.M. Fernandez-Mayoralas, N. Munoz-Jareno and B. Calleja-Perez, An open-label, prospective study of levitiracetam in children and adolescents with Tourette syndrome, *Eur J Paediatr Neurol* **13**(6) (2009), 541–545.
- [70] A.E. Cavanna, A. Schrag, D. Morley, M. Orth, M.M. Robertson, E. Joyce, H.D. Critchley and C. Selai, The Gilles de la Tourette Syndrome-Quality of Life Scale (GTS-QOL): development and validation, *Neurology* **71**(18) (2008), 1410–1416.
- [71] T. Pringsheim and T. Steeves, Pharmacological treatment for Attention Deficit Hyperactivity Disorder (ADHD) in children with comorbid tic disorders, *Cochrane Database of Systematic Reviews*, 2011, No. 4, CD007990.
- [72] P.R. Sanberg, R.D. Shytle and A.A. Silver, Treatment of Tourette's syndrome with mecamlamine in Tourette's syndrome: a two-year retrospective case study, *Lancet* **352**(9129) (1998), 705–706.
- [73] Y. Awaad, Tics in Tourette syndrome: New treatment options, *J Child Neurol* **14**(5) (1999), 316–319.
- [74] Y. Worbe, S. Palminteri, A. Hartmann, M. Vidailhet, S. Lehericy and M. Pessiglione, Reinforcement learning and gilles de la tourette syndrome: dissociation of clinical phenotypes and pharmacological treatments, *Arch Gen Psychiatry* **68**(12) (2011), 1257–1266.
- [75] P.R. Sanberg, A.A. Silver, R.D. Shytle, M.K. Philipp, D.W. Cahill, H.M. Fogelson and B.J. McConville, Nicotine for the treatment of Tourette's syndrome, *Pharmacol Ther* **74**(1) (1997), 21–25.
- [76] J.F. Lipinski, F.R. Sallee, C. Jackson and G. Sethuraman, Dopamine agonist treatment of Tourette disorder in children: results of an open-label trial of pergolide, *Mov Disord* **12**(3) (1997), 402–407.
- [77] B.J. McConville, P.R. Sanberg, M.H. Fogelson and J. King, The effects of nicotine plus haloperidol compared to nicotine only and placebo nicotine only in reducing tic severity and frequency in Tourette's disorder, *Biol Psychiatry* **31**(8) (1992), 832–840.
- [78] K.R. Muller-Vahl, The treatment of Tourette's syndrome: current opinions, *Expert Opin Pharmacother* **3**(7) (2002), 899–914.
- [79] A.A. Silver, R.D. Shytle and P.R. Sanberg, Mecamlamine in Tourette's syndrome: a two-year retrospective case study, *J Child Adolesc Psychopharmacol* **10**(2) (2000), 59–68.
- [80] P. Toren, N. Laor, D.J. Cohen, L. Wolmer and A. Weizman, Ondansetron treatment in patients with Tourette's syndrome, *Int Clin Psychopharmacol* **14**(6) (1999), 373–376.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

