

A NEW CHAOTIC ATTRACTOR FROM 2D DISCRETE MAPPING VIA BORDER-COLLISION PERIOD-DOUBLING SCENARIO

ZERAOULIA ELHADJ

Received 12 April 2005

The following map is studied: $(x, y) \rightarrow (1 + a(|x| - y^2) + y, bx)$. It is proved numerically that this model can display two different chaotic attractors, one is new and the other is a Lozi-type attractor. The new chaotic attractor is allowed via a border-collision period-doubling scenario, which is different from the classical period-doubling bifurcation.

1. Introduction

The discrete mathematical models are gotten directly via scientific experiences, or by the use of the Poincaré section for the study of a continuous model. One of these models is the Henon map. Many papers have described chaotic systems, one of the most famous being a two-dimensional discrete map which models the original Henon map [3, 4, 5, 7, 8]. Moreover, it is possible to change the form of the Henon map for obtaining others chaotic attractors [2, 6], this type of applications is used in secure communications using the notions of chaos.

The Lozi map is 2D noninvertible iterated map proposed by Lozi [6] as follow:

$$(x, y) \longrightarrow (1 - a|x| + y, bx). \quad (1.1)$$

This model gives a chaotic attractor called *Lozi attractor* and its shape is resembled to the one shown in Figure 2.1.

2. The proposed model

In this article, we essentially study the following modified Lozi map:

$$(x, y) \longrightarrow (1 + a(|x| - y^2) + y, bx). \quad (2.1)$$

We show numerically that the new model (2.1) can display two different chaotic attractors; including the classical Lozi-type attractor as shown in Figure 2.1 to Figure 4.2.




Figure 2.1. A Lozi-type chaotic attractor obtained from system (2.1) for $a = -1.8, b = 0.3$.




Figure 2.2. A Lozi-type chaotic attractor obtained from system (2.1) for $a = -1.8, b = 0.5$.

3. Comparison with the Lozi system

The system (2.1) has the same complexity as the Lozi system (1.1), they are both two-dimensional discrete noninvertible maps. However, the two models are topologically not equivalent because the Lozi system (1.1) is a piecewise-linear map, but model (2.1) is a nonlinear system. In addition; it can be rigorously proved that a non-singular homeomorphism that transforms each system to other does not exist. The proof needed some straightforward but tedious algebra which leads to a system of algebraic equations without solutions.

4. Route to chaos

It is well known that while varying the parameter a , the Hénon attractor is obtained via a period-doubling bifurcation route to chaos as a typical future [3], unless for Lozi map, no




Figure 4.1. A typical orbit of system (2.1) obtained for $a = 1.35$, $b = 0.3$.




Figure 4.2. The new chaotic attractor obtained for $a = 1.4$, $b = 0.3$.

period doubling route to chaos is allowed, and the attractor goes directly from a border-collision bifurcation developed from a stable periodic orbit [4], and for the same case the new chaotic attractor given by system (2.1) is obtained from a border-collision period-doubling bifurcation scenario [1]; this scenario (see Figure 4.3) is formed by a sequence of pairs of bifurcations, whereby each pair consists of a border-collision bifurcation and a pitchfork bifurcation. Thus, the three chaotic attractors go via different and distinguishable route to chaos as a typical future.

5. Conclusion

This paper reports the finding of a new two-dimensional discrete chaotic attractor obtained via direct modification of the Lozi mapping. The new chaotic attractor is allowed via a border-collision period-doubling scenario. More detailed analysis about the structure, dynamics and bifurcations of the new system (2.1) will be provided in near future.




Figure 4.3. Border-collision period-doubling scenario route to chaos observed for system (2.1).

References

- [1] V. Avrutin and M. Schanz, *Border-collision period-doubling scenario*, Phys. Rev. E (3) **70** (2004), no. 2, 026222, 11.
- [2] M. A. Aziz-Alaoui, C. Robert, and C. Grebogi, *Dynamics of a Hénon-Lozi-type map*, Chaos Solitons Fractals **12** (2001), no. 12, 2323–2341.
- [3] M. Benedicks and L. Carleson, *The dynamics of the Hénon map*, Ann. of Math. (2) **133** (1991), no. 1, 73–169.
- [4] Y. Cao and Z. Liu, *Strange attractors in the orientation-preserving Lozi map*, Chaos Solitons Fractals **9** (1998), no. 11, 1857–1863.
- [5] M. Hénon, *A two-dimensional mapping with a strange attractor*, Comm. Math. Phys. **50** (1976), no. 1, 69–77.
- [6] R. Lozi, *Un attracteur étrange du type attracteur de Hénon*, J. Phys. Colloque C5 **39** (1978), Supplément au no. 8, 9–10 (French).
- [7] F. R. Marotto, *Chaotic behavior in the Hénon mapping*, Comm. Math. Phys. **68** (1979), no. 2, 187–194.
- [8] M. Misiurewicz, *Strange attractors for the Lozi mappings*, Nonlinear Dynamics (Internat. Conf., New York, 1979) (R. G. Heman, ed.), Ann. New York Acad. Sci., vol. 357, New York Academy of Sciences, New York, 1980, pp. 348–358.

Zeraoula Elhadj: Department of Mathematics, University of Tébessa, 12000 Tébessa, Algeria
E-mail address: zelhadj12@yahoo.fr



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

