

## Research Article

# On $p$ -Adic Analogue of $q$ -Bernstein Polynomials and Related Integrals

T. Kim,<sup>1</sup> J. Choi,<sup>1</sup> Y. H. Kim,<sup>1</sup> and L. C. Jang<sup>2</sup>

<sup>1</sup> Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

<sup>2</sup> Department of Mathematics and Computer Science, KonKuk University, Chungju 380-701, Republic of Korea

Correspondence should be addressed to T. Kim, tkkim@kw.ac.kr

Received 17 September 2010; Accepted 22 December 2010

Academic Editor: Binggen Zhang

Copyright © 2010 T. Kim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Recently, Kim's work (in press) introduced  $q$ -Bernstein polynomials which are different Phillips'  $q$ -Bernstein polynomials introduced in the work by (Phillips, 1996; 1997). The purpose of this paper is to study some properties of several type Kim's  $q$ -Bernstein polynomials to express the  $p$ -adic  $q$ -integral of these polynomials on  $\mathbb{Z}_p$  associated with Carlitz's  $q$ -Bernoulli numbers and polynomials. Finally, we also derive some relations on the  $p$ -adic  $q$ -integral of the products of several type Kim's  $q$ -Bernstein polynomials and the powers of them on  $\mathbb{Z}_p$ .

## 1. Introduction

Let  $C[0, 1]$  denote the set of continuous functions on  $[0, 1]$ . For  $0 < q < 1$  and  $f \in C[0, 1]$ , Kim introduced the  $q$ -extension of Bernstein linear operator of order  $n$  for  $f$  as follows:

$$\mathbb{B}_{n,q}(f | x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} [x]_q^k [1-x]_{1/q}^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x, q), \quad (1.1)$$

where  $[x]_q = (1 - q^x)/(1 - q)$  (see [1]). Here  $\mathbb{B}_{n,q}(f | x)$  is called Kim's  $q$ -Bernstein operator of order  $n$  for  $f$ . For  $k, n \in \mathbb{Z}_+ (= \mathbb{N} \cup \{0\})$ ,  $B_{k,n}(x, q) = \binom{n}{k} [x]_q^k [1-x]_{1/q}^{n-k}$  are called the Kim's  $q$ -Bernstein polynomials of degree  $n$  (see [2-6]).

In [7], Carlitz defined a set of numbers  $\xi_k = \xi_k(q)$  inductively by

$$\xi_0 = 1, \quad (q\xi + 1)^k - \xi_k = \begin{cases} 1 & \text{if } k = 1, \\ 0 & \text{if } k > 1, \end{cases} \quad (1.2)$$

with the usual convention of replacing  $\xi^k$  by  $\xi_k$ . These numbers are  $q$ -analogues of ordinary Bernoulli numbers  $B_k$ , but they do not remain finite for  $q = 1$ . So he modified the definition as follows:

$$\beta_{0,q} = 1, \quad q(q\beta + 1)^k - \beta_{k,q} = \begin{cases} 1 & \text{if } k = 1, \\ 0 & \text{if } k > 1, \end{cases} \quad (1.3)$$

with the usual convention of replacing  $\beta^k$  by  $\beta_{k,q}$  (see [7]). These numbers  $\beta_{n,q}$  are called the  $n$ th Carlitz  $q$ -Bernoulli numbers. And Carlitz's  $q$ -Bernoulli polynomials are defined by

$$\beta_{k,q}(x) = (q^x\beta + [x]_q)^k = \sum_{i=0}^k \binom{k}{i} \beta_{i,q} q^{ix} [x]_q^{k-i}. \quad (1.4)$$

As  $q \rightarrow 1$ , we have  $\beta_{k,q} \rightarrow B_k$  and  $\beta_{k,q}(x) \rightarrow B_k(x)$ , where  $B_k$  and  $B_k(x)$  are the ordinary Bernoulli numbers and polynomials, respectively.

Let  $p$  be a fixed prime number. Throughout this paper,  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ , and  $\mathbb{C}_p$  will denote the ring of rational integers, the field of rational numbers, the ring of  $p$ -adic integers, the field of  $p$ -adic rational numbers and the completion of algebraic closure of  $\mathbb{Q}_p$ , respectively. Let  $\nu_p$  be the normalized exponential valuation of  $\mathbb{C}_p$  such that  $|p|_p = p^{-\nu_p(p)} = 1/p$ .

Let  $q$  be regarded as either a complex number  $q \in \mathbb{C}$  or a  $p$ -adic number  $q \in \mathbb{C}_p$ . If  $q \in \mathbb{C}$ , we assume  $|q| < 1$ , and if  $q \in \mathbb{C}_p$ , we normally assume  $|1 - q|_p < 1$ .

We say that  $f$  is a uniformly differentiable function at a point  $a \in \mathbb{Z}_p$  and denote this property by  $f \in \text{UD}(\mathbb{Z}_p)$  if the difference quotient  $F_f(x, y) = (f(x) - f(y))/(x - y)$  has a limit  $f'(a)$  as  $(x, y) \rightarrow (a, a)$  (see [1, 3, 8–13]).

For  $f \in \text{UD}(\mathbb{Z}_p)$ , let us begin with the expression

$$\frac{1}{[p^N]_q} \sum_{0 \leq x < p^N} q^x f(x) = \sum_{0 \leq x < p^N} f(x) \mu_q(x + p^N \mathbb{Z}_p), \quad (1.5)$$

representing a  $q$ -analogue of the Riemann sums for  $f$  (see [11]). The integral of  $f$  on  $\mathbb{Z}_p$  is defined as the limit as  $n \rightarrow \infty$  of the sums (if exists). The  $p$ -adic  $q$ -integral on a function  $f \in \text{UD}(\mathbb{Z}_p)$  is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \quad (1.6)$$

(see [11]).

As was shown in [3], Carlitz's  $q$ -Bernoulli numbers can be represented by  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  as follows:

$$\int_{\mathbb{Z}_p} [x]_q^m d\mu_q(x) = \beta_{m,q}, \quad \text{for } m \in \mathbb{Z}_+. \quad (1.7)$$

Also, Carlitz's  $q$ -Bernoulli polynomials  $\beta_{k,q}(x)$  can be represented

$$\beta_{m,q}(x) = \int_{\mathbb{Z}_p} [x+y]_q^m d\mu_q(y), \quad \text{for } m \in \mathbb{Z}_+, \quad (1.8)$$

(see [3]).

In this paper, we consider the  $p$ -adic analogue of Kim's  $q$ -Bernstein polynomials on  $\mathbb{Z}_p$  and give some properties of the several type Kim's  $q$ -Bernstein polynomials to represent the  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  of these polynomials. Finally, we derive some relations on the  $p$ -adic  $q$ -integral of the products of several type Kim's  $q$ -Bernstein polynomials and the powers of them on  $\mathbb{Z}_p$ .

## 2. $q$ -Bernstein Polynomials Associated with $p$ -Adic $q$ -Integral on $\mathbb{Z}_p$

In this section, we assume that  $q \in \mathbb{C}_p$  with  $|1-q|_p < 1$ .

From (1.5), (1.7) and (1.8), we note that

$$\begin{aligned} \int_{\mathbb{Z}_p} [1-x+x_1]_{1/q}^n d\mu_{1/q}(x_1) &= \frac{q^n}{(q-1)^{n-1}} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \frac{l+1}{q^{l+1}-1}, \\ \int_{\mathbb{Z}_p} [x+x_1]_q^n d\mu_q(x_1) &= \frac{1}{(1-q)^{n-1}} \sum_{l=0}^n \binom{n}{l} (-1)^l q^{lx} \frac{l+1}{1-q^{l+1}}. \end{aligned} \quad (2.1)$$

By (2.1), we get

$$(-1)^n q^n \int_{\mathbb{Z}_p} [x+x_1]_q^n d\mu_q(x_1) = \int_{\mathbb{Z}_p} [1-x+x_1]_{1/q}^n d\mu_{1/q}(x_1). \quad (2.2)$$

Therefore, we obtain the following theorem.

**Theorem 2.1.** For  $n \in \mathbb{Z}_+$ , one has

$$\int_{\mathbb{Z}_p} [1-x+x_1]_{1/q}^n d\mu_{1/q}(x_1) = (-1)^n q^n \int_{\mathbb{Z}_p} [x+x_1]_q^n d\mu_q(x_1). \quad (2.3)$$

By the definition of Carlitz's  $q$ -Bernoulli numbers and polynomials, we get

$$q^2 \beta_{n,q}(2) - (n+1)q^2 + q = q(q\beta + 1)^n = \beta_{n,q} \quad \text{if } n > 1. \quad (2.4)$$

Thus, we have the following proposition.

**Proposition 2.2.** *For  $n \in \mathbb{N}$  with  $n > 1$ , one has*

$$\beta_{n,q}(2) = \frac{1}{q^2} \beta_{n,q} + n + 1 - \frac{1}{q}. \quad (2.5)$$

It is easy to show that

$$[1-x]_{1/q}^n = (1-[x]_q)^n = (-1)^n q^n [x-1]_q^n. \quad (2.6)$$

Hence, we have

$$\int_{\mathbb{Z}_p} [1-x]_{1/q}^n d\mu_q(x) = (-1)^n q^n \int_{\mathbb{Z}_p} [x-1]_q^n d\mu_q(x). \quad (2.7)$$

By (1.8), we get

$$\int_{\mathbb{Z}_p} [1-x]_{1/q}^n d\mu_q(x) = (-1)^n q^n \beta_{n,q}(-1). \quad (2.8)$$

By Theorem 2.1 and (2.8), we see that

$$\int_{\mathbb{Z}_p} [1-x]_{1/q}^n d\mu_q(x) = (-1)^n q^n \beta_{n,q}(-1) = \beta_{n,1/q}(2). \quad (2.9)$$

From (2.9) and Proposition 2.2, we have

$$\int_{\mathbb{Z}_p} [1-x]_{1/q}^n d\mu_q(x) = \beta_{n,1/q}(2) = q^2 \beta_{n,1/q} + n + 1 - q. \quad (2.10)$$

By (1.7) and (2.10), we obtain the following theorem.

**Theorem 2.3.** *For  $n \in \mathbb{N}$  with  $n > 1$ , one has*

$$\int_{\mathbb{Z}_p} [1-x]_{1/q}^n d\mu_q(x) = q^2 \int_{\mathbb{Z}_p} [x]_{1/q}^n d\mu_{1/q}(x) + n + 1 - q. \quad (2.11)$$

Taking the  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  for one Kim's  $q$ -Bernstein polynomials, we get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x, q) d\mu_q(x) &= \binom{n}{k} \int_{\mathbb{Z}_p} [x]_q^k [1-x]_{1/q}^{n-k} d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \int_{\mathbb{Z}_p} [x]_q^{k+l} d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^{n-k} \binom{n-k}{l} (-1)^l \beta_{k+l,q}, \end{aligned} \tag{2.12}$$

and, by the  $q$ -symmetric property of  $B_{k,n}(x, q)$ , we see that

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x, q) d\mu_q(x) &= \int_{\mathbb{Z}_p} B_{n-k,n}\left(1-x, \frac{1}{q}\right) d\mu_q(x) \\ &= \binom{n}{k} \sum_{l=0}^k \binom{k}{l} (-1)^{k+l} \int_{\mathbb{Z}_p} [1-x]_{1/q}^{n-l} d\mu_q(x). \end{aligned} \tag{2.13}$$

For  $n > k + 1$ , by Theorem 2.3 and (2.13), one has

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x, q) d\mu_q(x) &= \binom{n}{k} \sum_{l=0}^k (-1)^{k+l} \binom{k}{l} \left[ n-l+1-q+q^2 \int_{\mathbb{Z}_p} [x]_{1/q}^{n-l} d\mu_{1/q}(x) \right] \\ &= \binom{n}{k} \sum_{l=0}^k (-1)^{k+l} \binom{k}{l} [n-l+1-q+q^2 \beta_{n-l,1/q}]. \end{aligned} \tag{2.14}$$

Let  $m, n, k \in \mathbb{Z}_+$  with  $m + n > 2k + 1$ . Then the  $p$ -adic  $q$ -integral for the multiplication of two Kim's  $q$ -Bernstein polynomials on  $\mathbb{Z}_p$  can be given by the following relation:

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x, q) B_{k,m}(x, q) d\mu_q(x) &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} [x]_q^{2k} [1-x]_{1/q}^{n+m-2k} d\mu_q(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} q \int_{\mathbb{Z}_p} [1-x]_{1/q}^{n+m-l} d\mu_q(x). \end{aligned} \tag{2.15}$$

By Theorem 2.3 and (2.15), we get

$$\begin{aligned}
& \int_{\mathbb{Z}_p} B_{k,n}(x,q)B_{k,m}(x,q)d\mu_q(x) \\
&= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \left[ n+m-l+1-q+q^2 \int_{\mathbb{Z}_p} [x]_{1/q}^{n+m-l} d\mu_{1/q}(x) \right] \\
&= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{2k} \binom{2k}{l} (-1)^{l+2k} \left[ n+m-l+1-q+q^2 \beta_{n+m-l,1/q} \right].
\end{aligned} \tag{2.16}$$

By the simple calculation, we easily get

$$\begin{aligned}
\int_{\mathbb{Z}_p} B_{k,n}(x,q)B_{k,m}(x,q)d\mu_q(x) &= \binom{n}{k} \binom{m}{k} \int_{\mathbb{Z}_p} [x]_q^{2k} [1-x]_{1/q}^{n+m-2k} d\mu_q(x) \\
&= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \int_{\mathbb{Z}_p} [x]_q^{l+2k} d\mu_q(x) \\
&= \binom{n}{k} \binom{m}{k} \sum_{l=0}^{n+m-2k} \binom{n+m-2k}{l} (-1)^l \beta_{l+2k,q}.
\end{aligned} \tag{2.17}$$

Continuing this process, we obtain

$$\begin{aligned}
\int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k,n_i}(x,q) \right) d\mu_q(x) &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \int_{\mathbb{Z}_p} [x]_q^{sk} [1-x]_{1/q}^{n_1+\dots+n_s-sk} d\mu_q(x) \\
&= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \int_{\mathbb{Z}_p} [1-x]_{1/q}^{n_1+\dots+n_s-l} d\mu_q(x).
\end{aligned} \tag{2.18}$$

Let  $s \in \mathbb{N}$  and  $n_1, \dots, n_s, k \in \mathbb{Z}_+$  with  $n_1 + n_2 + \dots + n_s > sk + 1$ . By Theorem 2.3 and (2.18), we get

$$\begin{aligned}
& \int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k,n_i}(x,q) \right) d\mu_q(x) \\
&= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \left\{ \sum_{i=1}^s n_i - l + 1 - q + q^2 \int_{\mathbb{Z}_p} [x]_{1/q}^{n_1+\dots+n_s-l} d\mu_{1/q}(x) \right\} \\
&= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \left\{ \sum_{i=1}^s n_i - l + 1 - q + q^2 \beta_{n_1+\dots+n_s-l,1/q} \right\}.
\end{aligned} \tag{2.19}$$

From the definition of binomial coefficient, we note that

$$\begin{aligned}
 & \int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k,n_i}(x, q) \right) d\mu_q(x) \\
 &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \int_{\mathbb{Z}_p} [x]_q^{sk} [1-x]_{1/q}^{n_1+\dots+n_s-sk} d\mu_q(x) \\
 &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \int_{\mathbb{Z}_p} [x]_q^{sk+l} d\mu_q(x) \\
 &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \beta_{sk+l,q},
 \end{aligned} \tag{2.20}$$

where  $s \in \mathbb{N}$  and  $n_1, \dots, n_s, k \in \mathbb{Z}_+$ .

By (2.19) and (2.20), we obtain the following theorem.

**Theorem 2.4.** (I) For  $s \in \mathbb{N}$  and  $n_1, \dots, n_s, k \in \mathbb{N}$  with  $n_1 + n_2 + \dots + n_s > sk + 1$ , one has

$$\begin{aligned}
 & \int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k,n_i}(x, q) \right) d\mu_q(x) \\
 &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \left\{ \sum_{i=1}^s n_i - l + 1 - q + q^2 \beta_{n_1+\dots+n_s-l,1/q} \right\}.
 \end{aligned} \tag{2.21}$$

(II) For  $s \in \mathbb{N}$  and  $n_1, \dots, n_s, k \in \mathbb{Z}_+$ , one has

$$\int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k,n_i}(x, q) \right) d\mu_q(x) = \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \beta_{sk+l,q}. \tag{2.22}$$

By Theorem 2.4, we obtain the following corollary.

**Corollary 2.5.** For  $s \in \mathbb{N}$  and  $n_1, \dots, n_s, k \in \mathbb{N}$  with  $n_1 + n_2 + \dots + n_s > sk + 1$ , one has

$$\begin{aligned}
 & \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{sk+l} \left\{ \sum_{i=1}^s n_i - l + 1 - q + q^2 \beta_{n_1+\dots+n_s-l,1/q} \right\} \\
 &= \sum_{l=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{l} (-1)^l \beta_{sk+l,q}.
 \end{aligned} \tag{2.23}$$

Let  $s \in \mathbb{N}$  and  $m_1, \dots, m_s, n_1, \dots, n_s, k \in \mathbb{Z}_+$  with  $m_1 n_1 + \dots + m_s n_s > (m_1 + \dots + m_s) k + 1$ . Then one has

$$\begin{aligned}
\int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k, n_i}^{m_i}(x, q) \right) d\mu_q(x) &= \left( \prod_{i=1}^s \binom{n_i}{k}^{m_i} \right) \sum_{l=0}^{k \sum_{i=1}^s m_i} \binom{k \sum_{i=1}^s m_i}{l} (-1)^{k \sum_{i=1}^s m_i - l} \\
&\quad \times \int_{\mathbb{Z}_p} [1-x]_q^{\sum_{i=1}^s n_i m_i - l} d\mu_q(x) \\
&= \left( \prod_{i=1}^s \binom{n_i}{k}^{m_i} \right) \sum_{l=0}^{k \sum_{i=1}^s m_i} \binom{k \sum_{i=1}^s m_i}{l} (-1)^{k \sum_{i=1}^s m_i - l} \\
&\quad \times \left\{ \left( \sum_{i=1}^s m_i n_i - l + 1 \right) - q + q^2 \int_{\mathbb{Z}_p} [x]_{1/q}^{\sum_{i=1}^s n_i m_i - l} d\mu_{1/q}(x) \right\} \\
&= \left( \prod_{i=1}^s \binom{n_i}{k}^{m_i} \right) \sum_{l=0}^{k \sum_{i=1}^s m_i} \binom{k \sum_{i=1}^s m_i}{l} (-1)^{k \sum_{i=1}^s m_i - l} \\
&\quad \times \left\{ \left( \sum_{i=1}^s m_i n_i - l + 1 \right) - q + q^2 \beta_{n_1 m_1 + \dots + n_s m_s - l, 1/q} \right\}. \tag{2.24}
\end{aligned}$$

From the definition of binomial coefficient, one has

$$\begin{aligned}
&\int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k, n_i}^{m_i}(x, q) \right) d\mu_q(x) \\
&= \left( \prod_{i=1}^s \binom{n_i}{k}^{m_i} \right) \sum_{l=0}^{\sum_{i=1}^s n_i m_i - k \sum_{i=1}^s m_i} \binom{\sum_{i=1}^s n_i m_i - k \sum_{i=1}^s m_i}{l} (-1)^l \\
&\quad \times \int_{\mathbb{Z}_p} [x]_q^{(m_1 + \dots + m_s)k + l} d\mu_q(x) \tag{2.25} \\
&= \left( \prod_{i=1}^s \binom{n_i}{k}^{m_i} \right) \sum_{l=0}^{\sum_{i=1}^s n_i m_i - k \sum_{i=1}^s m_i} \binom{\sum_{i=1}^s n_i m_i - k \sum_{i=1}^s m_i}{l} \\
&\quad \times (-1)^l \beta_{(m_1 + \dots + m_s)k + l, q}.
\end{aligned}$$

By (2.24) and (2.25), we obtain the following theorem.



**Theorem 2.6.** For  $s \in \mathbb{N}$  and  $m_1, \dots, m_s, n_1, \dots, n_s, k \in \mathbb{Z}_+$  with  $m_1 n_1 + \dots + m_s n_s > (m_1 + \dots + m_s)k + 1$ , one has

$$\begin{aligned} & \sum_{l=0}^{k \sum_{i=1}^s m_i} \binom{k \sum_{i=1}^s m_i}{l} (-1)^{k \sum_{i=1}^s m_i - l} \left\{ \binom{\sum_{i=1}^s m_i n_i - l + 1}{l} - q + q^2 \beta_{n_1 m_1 + \dots + n_s m_s - l, 1/q} \right\} \\ &= \sum_{l=0}^{\sum_{i=1}^s n_i m_i - k \sum_{i=1}^s m_i} \binom{\sum_{i=1}^s n_i m_i - k \sum_{i=1}^s m_i}{l} (-1)^l \beta_{(m_1 + \dots + m_s)k + l, q}. \end{aligned} \quad (2.26)$$

## Acknowledgment

This paper was supported by the research grant of Kwangwoon University in 2010.

## References

- [1] T. Kim, "A note on  $q$ -Bernstein polynomials," *Russian Journal of Mathematical Physics*. In press.
- [2] M. Acikgoz and S. Araci, "A study on the integral of the product of several type Bernstein polynomials," *IST Transaction of Applied Mathematics-Modelling and Simulation*, vol. 1, no. 1, pp. 10–14, 2010.
- [3] T. Kim, "On a  $q$ -analogue of the  $p$ -adic log gamma functions and related integrals," *Journal of Number Theory*, vol. 76, no. 2, pp. 320–329, 1999.
- [4] G. M. Phillips, "On generalized Bernstein polynomials," in *Numerical Analysis*, pp. 263–269, World Scientific, River Edge, NJ, USA, 1996.
- [5] G. M. Phillips, "Bernstein polynomials based on the  $q$ -integers," *Annals of Numerical Analysis*, vol. 4, pp. 511–514, 1997.
- [6] Y. Simsek and M. Acikgoz, "A new generating function of ( $q$ -) Bernstein-type polynomials and their interpolation function," *Abstract and Applied Analysis*, vol. 2010, Article ID 769095, 12 pages, 2010.
- [7] L. Carlitz, " $q$ -Bernoulli numbers and polynomials," *Duke Mathematical Journal*, vol. 15, pp. 987–1000, 1948.
- [8] M. Cenkci, V. Kurt, S. H. Rim, and Y. Simsek, "On ( $i, q$ ) Bernoulli and Euler numbers," *Applied Mathematics Letters*, vol. 21, no. 7, pp. 706–711, 2008.
- [9] L.-C. Jang, "A new  $q$ -analogue of Bernoulli polynomials associated with  $p$ -adic  $q$ -integrals," *Abstract and Applied Analysis*, vol. 2008, Article ID 295307, 6 pages, 2008.
- [10] T. Kim, "Barnes-type multiple  $q$ -zeta functions and  $q$ -Euler polynomials," *Journal of Physics A*, vol. 43, no. 25, Article ID 255201, 11 pages, 2010.
- [11] T. Kim, " $q$ -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [12] T. Kim, L.-C. Jang, and H. Yi, "A note on the modified  $q$ -Bernstein polynomials," *Discrete Dynamics in Nature and Society*, vol. 2010, Article ID 706483, 12 pages, 2010.
- [13] B. A. Kupershmidt, "Reflection symmetries of  $q$ -Bernoulli polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 12, supplement 1, pp. 412–422, 2005.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

