

distal joints darker, femora with a dark anterior lateral line, much fainter on the posterior pairs, all joints with rows of small hairs, no spines, tarsal comb of 6 or 7 curved bristles on the fourth metatarsus; *palpus*, large for the size of the spider, shorter than the cephalothorax, femur pale and bent, other joints dark, patella short and much swollen on the dorsal side, tibia very small and pressed close to the cymbium, palpal organ short, barrel-shaped, embolus probably a short black spine at the tip.

Holotype: ♂ Texas; Dallas, on the outside of a house, 9 July 1936, (Jones); in MCZ.

A CORRECTION.—In *Pysche*, Vol. 56, No. 1, I published a synonymic list involving some species in *Pseudomyrma* and other genera of ants (pp. 41–49). On page 43 occur two errors, the first of which was made at the printing office *after* the page proof had been read by both author and editor. Page 43, line 5, should read “*Pseudomyrma spinicola* subsp. *infernalis*”, not “subsp. *sclerosa*.” Line 9 on the same page mistakenly omits an “e” from “subsp. *scelerosa*”, the latter being Wheeler’s original spelling. Enzmann’s transcription of the name was “*sclerosa*”.—W. L. BROWN, JR., Biological Laboratories, Harvard University.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

