TABANIDAE OF NEW HAMPSHIRE

BY R. L. BLICKLE¹

University of New Hampshire

Many species of Tabanidae are of economic importance in the state as pests of man and of livestock. In general, the species of the genus Tabanus are more important as regards livestock, while those of the genus Chrysops are irritating to man. The following horseflies, listed in order of their appearance during the summer, are of economic importance; Tabanus lasiopthalmus, T. typhus, T. lineola, T. trispilus, T. aurilimbus, and T. sulcifrons. Along the coast T. nigrovittatus in mid-summer and T. actaeon in late summer have been numerous enough at times to be important. Other species present at the same time as those mentioned above, though not numerous, contribute to the losses attributed to horseflies. The deerflies, although very numerous, do not irritate the cattle or cause as much loss of blood as do the larger horseflies. The deerflies are important as pests of man in recreational areas and in tracts where lumbering operations are being conducted. Those species causing discomfort to man because of their abundance are: Chrysops mitis, C. sordida, C. carbonaria, and C. excitans in early summer. C. lateralis, C. striata, C. *univittata*, and C. vittata are important along the seashore and causes more discomfort to many than any other species of Tabanidae.

The list of species and subspecies is based mainly on material in the University of New Hampshire collection. Incorporated in the collection are the specimens donated by the late S. A. Shaw. Mr. Shaw was an active collector of insects, especially Diptera, from 1890 until 1934. Other collectors contributing to the collection have been: R. L. Blickle, J. G. Conklin, W. F. Fiske, A. G. Humes, P. R. Lowry, W. J. Morse, H. W. Smith and C. M. Weed. Additional records were obtained from the Museum of Com-

¹Published with the permission of the Director as Scientific Contribution No. 156 of the New Hampshire Agricultural Experiment Station. parative Zoology and from the Boston Society collection, now at Boston University. The examination of these collections was made possible through the courtesy of Drs. J. Bequaert and A. G. Humes, respectively.

Published lists recording species taken in New Hampshire are: Johnson (1925, 1925a), Brennan (1935), Stone (1938), Philip (1947, 1950), and Fairchild (1950). In all forty-nine species and subspecies are recorded in the above works. The present list contains seventy-one species and subspecies. Four species recorded from both Maine and Massachusetts are also listed. Judging from the distribution of tabanids in the neighboring states and in Canada, the number recorded will eventually be approximately ninety.

Locality records are given as townships. Only the extreme dates of capture are listed. Most of the records are from the southeastern part of the state and from the White Mountain area. The western part of the state north of the Mt. Monadnock area is poorly represented.

The author is indebted to Dr. L. L. Pechuman for the determination of the flies.

Stonemyia Brennan

rasa (Loew): Durham, Gilmanton, Hampton, Lee, Mt. Washington, New Durham, Pelham, July 19 to Aug. 29.

tranquilla (Osten Sacken): Crawford Notch, Durham, Gilmanton, Hampton, Kearsarge, Lee, New Durham, Mt. Washington, North Conway, Ossipee, Pinkham Notch, Twin Mts., White Mountains (type loc.) July 18 to Aug. 23.

Chrysops Meigen

aberrans Philip: Alton, Durham, Gilmanton, Lee, Madbury, New Durham. June 15 to Sept. 9.

- amazon Daecke: Durham, Lee, Rochester. June 15 to July 13,
- atlantica Pechuman: Hampton (S. A. Shaw). July 1 to Aug. 21.

callida Osten Sacken: Barrington, Durham, Gilmanton, Hampton, Lee, Milford. June 12 to Aug. 4.

carbonaria Walker: Bretton Woods, Colebrook, Conway,

Durham, Gorham, Hampton, Hanover, Intervale, Lee, Madbury, Merrimack, Ossipee, Pittsburg, Twin Mts. May 7 to July 14.

celer Osten Sacken: Colebrook, Dummer, Durham, Hampton, Lee, Milford, Pittsburg, Rye, June 1 to July 23.

cuclux Whitney: Bretton Woods, Colebrook, Durham, Hampton, Madbury, Milford (type loc.), Moultonboro, Pittsburg, White Mts., May 10 to July 14.

cursim Whitney: Milford (type loc.). July.

delicatula Osten Sacken: Alton, Durham, Gilmanton, Lee, Madbury, Madison, New Durham, North Conway (type loc.), Webster. July 12 to Aug. 14.

dimmocki Hine: Lee (R. L. Blickle). July 8.

excitans Walker: Antrim, Belmont, Bretton Woods, Colebrook, Dixville, Durham, Dummer, Gorham, Lee, Madbury, Newfields, North Conway, Ossipee, Pittsburg, Rye, Tamworth. June 3 to July 12.

frigida Osten Sacken: Durham, Lee, Milford, New Durham, North Conway, White Mts. June 11 to July 14. frigida xantha Philip: Milford, Franconia. July 8.

- fuliginosa Wiedemann: Durham, Hampton, Lee. June 6 to July 2.
- geminata Wiedemann: Belknap, Belmont, Durham, Greenfield, Hill, Lee, Lyme, Madbury, Milford, New Durham, Wilton. July 5 to July 25.
- geminata impunctata Krober: Durham (R. L. Blickle). July 18.

inda Östen Sacken: Milford, Pittsburg. June 20 to July 1. *lateralis* Wiedemann: Barrington, Belmont, Bretton

Woods, Dublin, Durham, Gilford, Gilmanton, Hampton, Hanover, Intervale, Lee, Jaffrey, Madbury, Milford, Mt. Monadnock, New Durham, North Conway, Portsmouth, Rochester, Squam Lake, Mt. Washington, Woodstock. June 11 to Aug. 24.

mitis Osten Sacken: Bretton Woods, Dixville, Franconia, Jaffrey, Mt. Washington, Pittsburg, June 11 to July 7.

moecha Osten Sacken: Alton, Durham, Hampton, Lee, Madbury. June 1 to Aug. 24.

montana Osten Sacken: Alton, Durham, Lee. July 6 to July 18.

nigra Macquart: Alton, Belmont, Dixville, Durham,

Hampton, Madbury, Milford, Mt. Washington, Newfields, Pittsburg, Rochester. June 1 to Aug. 11.

nigribimbo Whitney: Milford (type loc.). July 9 to July 21.

pudica Osten Sacken: Durham, Lee (R. L. Blickle). July 4.

- sackeni Hine: Durham, Hampton, Lee, Milford, Newfields. June 17 to Aug. 8.
- shermani Hine: Berlin, Dixville (R. L. Blickle). June 19 to June 20.
- sordida Osten Sacken: Bretton Woods, Colebrook, Dixville, Dummer, Franconia, Mt. Washington, Pittsburg, Twin Mts. (type loc.). June 19 to July 16.
- striata Osten Sacken: Belmont, Durham, Gilmanton, Hampton, Lee, Madbury, New Durham, Sanbornton. June 22 to July 30.
- univittata Macquart: Belmont, Durham, Gilmanton, Hampton, Lee, Madbury, Milford, New Durham, Newfields, Ossipee. May 21 to Sept. 11.
- vittata Wiedemann: Belmont, Durham, Gilmanton, Greenfield, Hampton, Lee, Madbury, Milford, New Durham, Pelham, Rye, Sanbornton. June 19 to Sept. 9.
- wiedemanni Krober: Alton, Durham, Epping, Hampton, Lee, Madbury, Milford, New Durham. June 12 to Sept. 9.

Atylotus Osten Sacken

bicolor (Wiedemann): Hampton, Webster. July 15 to Aug. 5.

pemeticus (Johnson): Recorded from Maine and Massachusetts (Philip 1947).

thoracicus (Hine): Recorded from Maine and Massachusetts (Philip 1947).

Tabanus L. (Hybomitra Enderlein)

- affinis Kirby: Bretton Woods, Intervale, Jaffrey, Mt. Monadnock, New Durham, Peterboro, Shelburne, Mt. Washington. June 21 to July 11.
- aurilimbus Stone: Belmont, Durham, Gilmanton, Hampton, Epping, Madbury, Milford, New Durham, North Hampton. June 29 to Aug. 5.

- astutus Osten Sacken: Pinkham Notch, White Mts. (type loc.). Aug. 21-22.
- brennani Stone: Franconia (type loc.). July 14.
- *cinctus* Fabricius: Dublin, Durham, Lee, Madbury, Union. July 23 to Aug. 2.
- epistates Osten Sacken: Durham, Lee, Mt. Monadnock, Shelburne. July 4 to July 25.
- hinei Johnson: Lee, Madbury (R. L. Blickle). June 28 to July 25.
- *illotus* Osten Sacken: Pittsburg (Morse, Blickle). June 20.
- lasiopthalmus Macquart: Bretton Woods, Dummer, Durham, Hampton, Hanover, Jackson, Jefferson, Lee, Newfields, Pelham, Pittsburg, Rye, Twin Mts. May 30 to Aug. 15.
- longiglossus Philip: Recorded from Maine and Massachusetts (Philip 1947).
- metabolus McDunnough: Hampton, Jaffrey, Milford, Passaconaway, Pittsburg.

microcephalus Osten Sacken: Lee, Jackson, Madbury, New Durham, Pinkham Notch, Rumney, Twin Mts., White Mts. (type loc.). July 8 to Sept. 9.

- minusculus Hine: Philip (1950).
- nudus McDunnough: Durham, Hampton, Jackson, Jaffrey, Lee, New Durham, Passaconaway, Pittsburg. May 3 to June 20.
- septentrionalis Loew: Recorded from Maine and Massachusetts (Philip 1947).
- trepidus McDunnough: Colebrook, Dummer, Durham, Hampton, Lee. June 10 to Aug. 10.
- trispilus Wiedemann: Barrington, Dublin, Durham, Epping, Franconia, Hampton, Lee, Madbury, Newfields, Pittsfield. July 4 to Aug. 8.
- typhus Whitney: Barrington, Durham, Freedom, Lee, Madbury, Milford (type loc.), New Durham, North Conway. June 15 to July 27.
- *zonalis* Kirby: Lee, Milford, Passaconaway. June 8 to July 15.

Tabanus L. (Tabanus s.str.)

- actaeon Osten Sacken: Amherst, Dublin, Durham, Hampton, Lee, Milford, Pelham, Rye. Aug. 9 to Sept. 17.
- americanus Forster: Dublin. (Johnson 1925 a). July.
- atratus Fabricius: Durham, Hampton, Lee, Pelham. June 30 to Aug. 28.
- atratus fulvopilosus Johnson: Durham (H. W. Smith). July 22.
- atratus nantuckensis Hine: Hampton, Newfields, Rochester. July 9 to Sept. 13.
- catenatus Walker: Durham, Hampton, Jaffrey, Lee, Rye. July 7 to Aug. 27.
- fairchildi Stone: Alstead, Hillsboro. July 1 to July 16.
- *lineola* Fabricius: Barrington, Durham, Gilmanton, Hampton, Lee, Madbury.
- *lineola scutellaris* Walker: Durham, Hampton, Lee. June 17 to July 20.
- *nigripes* Wiedemann: Hampton, Milford. July 3 to July 14.
- nigrovittatus Macquart: Durham, Hampton, Newmarket, Portsmouth, Rye. June 30 to Aug. 8.
- nivosus Osten Sacken: Durham, Freedom, Hampton, Lee, New Durham. June 12 to July 31.
- *pumilus* Macquart: Barrington, Durham, Lee, Madbury, Portsmouth. July 1 to July 31.
- quinquevittatus Wiedemann: Durham, Gilmanton, Lee, Madbury, New Durham, Newfields, Newmarket. July 2 to Sept. 21.
- reinwardtii Wiedemann: Durham, Newfields. July 11 to Aug. 1.
- sparus Whitney: Durham, Hampton, Lee, Milford (type loc.), Newfields, Rye. June 10 to July 20.
- stygius Say: Durham. (no date).
- *sulcifrons* Macquart: Durham, Hampton, Lee, Madbury. July 11 to Sept. 9.
- superjumentarius Whitney: Durham, Epping, Hampton, Lee, Madbury, Milford (type loc.). June 14 to Aug. 3.
- vivax Osten Sacken: Durham, Farmington, Gilmanton, Hampton, Lee, Strafford. July 4 to July 23.

LITERATURE CITED

BRENNAN, J. M.

1935. The Pangoniinae of Nearctic America. Diptera. Tabanidae. Bull. Univ. Kansas, 36:1-401.

FAIRCHILD, G. B.

1950. The Diptera or true flies of Connecticut. Family Tabanidae. Fasc. 4. Conn. State Geol. Nat. Hist. Surv. Bull., 75:3-31.

JOHNSON, C. W.

- 1925a. Fauna of New England 15. List of the Diptera or two-winged flies. Occas. Pap. Bost. Soc. Nat. Hist., VII: 1-326.
 - 1925b. Diptera of the Harris Collection. Proc. Bost. Soc. Nat. Hist., 38:57-90.

Philip, C. B.

- 1947. A catalog of the Blood-sucking Fly Family Tabanidae of the Nearctic Region North of Mexico. Amer. Mid. Nat., 37(2): 257-324.
- 1950. Corrections and Addenda to a Catalog of Nearctic Tabanidae. Amer. Mid. Nat., 43(2):430.

STONE, ALAN

1938. The Horseflies of the Subfamily Tabanidae of the Nearctic Region. U.S.D.A. Misc. Pub. No. 305: 1-168.



BioMed Research International

Zoology





Hindawi

Submit your manuscripts at http://www.hindawi.com





International Journal of Genomics





The Scientific World Journal



Journal of Signal Transduction

Genetics Research International



Anatomy Research International



International Journal of Microbiology



Biochemistry Research International



Advances in Bioinformatics



Enzyme Research



International Journal of Evolutionary Biology



Molecular Biology International



Journal of Marine Biology