

THE MALE GENITALIA OF BLATTARIA.
XI. PERISPHAERIINAE.*

BY LOUIS M. ROTH

Pioneering Research Laboratory

U. S. Army Natick Laboratories

Natick, Massachusetts 01760

In this paper I shall illustrate and characterize the male genitalia of 11 genera (indicated with an * in Table 1) of cockroaches which belong to the Perisphaeriinae. Three recent classifications which include these genera are shown in the table. I follow McKittrick (1964) in placing all of those discussed here under the Blaberidae: Perisphaeriinae, and Beier (1961) in using tribes to separate them. Rehn (1922) discussed the characteristics of *Pilema*, *Cyrtotria*, *Dero-calymma*, *Zuluia*, and *Hostilia*, and Princis (1963) devised keys to some South African species of the above genera, as well as those of *Bantua*, *Perisphaeria*, and *Blepharodera*.

MATERIALS AND METHODS

All genitalia were treated with 10% KOH, dehydrated, cleared, and mounted in Permount. The source of each of the specimens examined is given using the following abbreviations: (ANSP) = Academy of Natural Sciences, Philadelphia; (BMNH) = British Museum (Natural History), London; (CUZM) = Zoological Museum, Copenhagen; (L) = Zoological Institute, Lund, Sweden; (HO) = Hope Department of Entomology, Oxford, England; (MCZ) = Museum of Comparative Zoology, Harvard University, Cambridge, Mass. Locality data and names of specialists who identified the specimens, if known, follow these abbreviations; most of the species were determined by Princis or Rehn. The number preceding the abbreviations refers to the number assigned the specimen and its corresponding genitalia (on a slide) which are deposited in their respective museums.

RESULTS AND DISCUSSION

The normal arrangement of the phallomeres of male genitalia of the Perisphaeriinae, as seen in dorsal view, is illustrated in figure 1. Based on male genitalia, I suggest the erection of 4 tribes for the

*Manuscript received by the editor October 10, 1973

Table 1. Three recent classifications of some of the species of Blattaria discussed in this paper (marked with an *)

Princis (1960)	Beier (1961)	McKittrick (1964)
Derocalymmidae	Blaberidae	Blaberidae
Apterinae	Perisphaeriinae	Perisphaeriinae
<i>Aptera</i> Sauss.	<i>Apterini</i>	<i>Perisphaeria</i>
<i>Elliptica</i>	<i>Aptera</i>	(= <i>Hyposphaeria</i>)
Sauss. & Zehn.	<i>Elliptica</i>	<i>Derocalymma</i>
<i>Elliptoblatta</i>	<i>Elliptoblatta</i>	<i>Eustegasta</i> Gerst. ^a
Sauss. & Zehn.	<i>Gymnonyx</i>	<i>Gyna</i>
<i>Gymnonyx</i>	<i>Thoracopygia</i>	<i>Pilema</i>
Sauss. & Zehn.		<i>Perisphaerus</i>
<i>Thoracopygia</i>	Derocalymmini	<i>Isoniscus</i> Borg. ^a
Sauss. & Zehn.	<i>Perisphaerus</i>	
	<i>Glomeriblatta</i>	
Derocalymminae	<i>Pseudoglomeris</i>	
* <i>Bantua</i> Shelf.	<i>Pronaonota</i>	
<i>Compsagis</i>	<i>Cyrtotria</i>	
Chop.	<i>Hostilia</i>	
* <i>Cyrtotria</i>	<i>Compsagis</i>	
Stål	<i>Derocalymma</i>	
* <i>Derocalymma</i>	<i>Platysilpha</i>	
Burm.	and others	
<i>Glomeriblatta</i>		
Bey-Bienko	Perisphaeriini	
* <i>Hostilia</i>	<i>Perisphaeria</i>	
Stål	<i>Blepharodera</i>	
* <i>Perisphaerus</i>	<i>Poeciloblatta</i>	
Serv.	<i>Parasphaeria</i>	
* <i>Pilema</i> Sauss.	<i>Gyna</i>	
<i>Platysilpha</i>	<i>Paranauphoeta</i>	
Shelf.	<i>Paraplecta</i>	
<i>Pronaonota</i>	and others	
Sauss. & Zehn.		
* <i>Pseudoglomeris</i>	Brachycolini ^c	
Brunner	<i>Brachycola</i> Serv.	
* <i>Zuluia</i> Rehn	<i>Hormetica</i> Burm.	
	<i>Parahormetica</i>	
	Brunner	
	and others	

Princis (1960)	Beier (1961)	McKittrick (1964)
Perisphaeriidae		
Gyninae		
<i>Evea</i> Shelf.		
<i>Gyna</i> Brunner ^b		
<i>Paranauphoeta</i>		
Brunner		
<i>Paraplecta</i> Shelf.		
<i>Progonogamia</i>		
Rehn		
<i>Thliptoblatta</i>		
Sauss. & Zehn.		
Perisphaeriinae		
* <i>Blepharodera</i>		
Burm.		
<i>Parasphaeria</i>		
Brunner ^d		
* <i>Perisphaeria</i>		
Burm.		
<i>Poeciloblatta</i>		
Sauss. & Zehn.		
* <i>Trichoblatta</i>		
Sauss. & Zehn.		
Zetoborinae ^e		
<i>Lanxoblatta</i> Hebard		
<i>Phortioeca</i> Sauss.		
<i>Schistopeltis</i> Rehn		
<i>Schizopilia</i> Burm.		
<i>Tribonoidea</i> Shelf.		
<i>Tribonium</i> Sauss.		
<i>Zetobora</i> Burm.		
<i>Zetoborella</i> Hebard		

^aPrincis (1960) placed these 2 genera in Blaberidae: Laxtinae. Beier (1961) placed *Eustegasta* in the Panchlorinae: Laxtini. The ♂ genitalia of Panchlorinae are characterized by a marked reduction or absence of one or more of the phallomeres (Roth, 1971; Gurney and Roth, 1972). The genitalia of *Eustegasta* are quite different from members of the Panchlorinae (unpublished observations).

^bThe genitalia of *Gyna* spp. have been described (Roth, 1972).

^cI (1970a) followed McKittrick (1964) and placed this tribe in Blaberinae.

^dThe genitalia of *Parasphaeria ovata* (Blanchard) (South America) is more typical of members of the Epilamprinae and I placed it in this subfamily (Roth, 1973).

^eI (1970) followed McKittrick (1964) in placing this subfamily closer to the Blaberinae (Blaberoid Complex) than to the Perisphaeriinae (Epilamprid Complex).





Figure 1. (62 BMNH). Male genitalia (dorsal) of *Cyrtotria latipennis* (Kirby), British East Africa (from ♂ shown in Fig. 37). Left (L) and right (R) phallomeres numbered according to McKittrick (1964); C = cleft of L1; UL and LL — upper and lower lobes of L1, respectively.


genera studied here; additional tribes will undoubtedly be required for other genera of this subfamily (unpublished observations).

Perisphaeriini. — The upper lobe of L1 is usually about the same height or slightly larger than the lower lobe and it has a snoutlike extension (accidentally lost in some of the illustrated preparations); setae are absent from the lower lobe. The L2d is variable in shape, its surface distinctly spicular (at least in part), or reticulate (at least in part) and usually with a smooth lateral extension on the right side. R2 with or without a subapical incision. In this tribe I include the African genera *Perisphaeria*, *Blepharodera*, *Pilema*, *Hostilia*, *Bantua*, and *Cyrtotria*, which can be further subdivided into groups.


Tribal Group I. The entire or at least part of the surface of L2d is spicular (not reticulate): *Perisphaeria*, *Blepharodera*, *Pilema*, and *Hostilia*.




Figures 2-9. Adult males of *Perisphaeria* spp. 2. (1424 L). *P. virescens* (Brunner). (det. Princis). 3. (40 CUZM). *P. scabra* (Thunb.). Cap bono Spei, Dec. 1816. 4. (1420 L). *P. scabra*. Holub, South Africa (det. Princis). 5. (1419 L). *P. saxicola* (Esch.). Capetown, Table Mountain, 2400'. (det. Princis). 6. (1422 L). *P. ruficornis* (Sauss. and Zehn.). Swartbergpas, Platberg, Cape Province, South Africa, ca. 5000' (det. Princis). 7. (1421 L). *P. stylifera* Burm. Manhica (det. Princis). 8. (161 ANSP). *P. stylifera*. Durban. (det. by Rehn as *Hyposphaeria giebeliana* (Sauss. and Zehn.)). 9. (1423 L). *P. aspera* (Walker). Baberton, Levetzow (det. Princis). (scale = 5 mm).




Figures 10-18. Male genital phallomeres of *Perisphaeria* spp. 10-12. (1424 L). *P. virescens* (from ♂ shown in Fig. 2). 13-15. (40 CUZM). *P. scabra* (from ♂ shown in Fig. 3). 16-18. (1420 L). *P. scabra* (from ♂ shown in Fig. 4). (scale = 0.2 mm).




Figures 19-27. Male genital phallomeres of *Perisphaeria* and *Cyrtotria* spp. 19-21. (1419 L). *P. saxicola* (from ♂ shown in Fig. 5). 22-24. (1422 L). *P. ruficornis* (from ♂ shown in Fig. 6). 25-27. (15 HO). *Cyrtotria* sp. (from ♂ shown in Fig. 105). (scale = 0.2 mm).




Figures 28-36. Male genital phallomeres of *Perisphaeria* spp. 28-30. (1421 L). *P. stylifera* (from ♂ shown in Fig. 7). 31-33. (161 ANSP). *P. stylifera* (from ♂ shown in Fig. 8). 34-36. (1423 L). *P. aspera* (from ♂ shown in Fig. 9). (scale = 0.2 mm).



Figures 37-42. Adult males of *Cyrtotria*, *Pilema*, and *Hostilia* spp. 37. (62 BMNH). *C. latipennis* (holotype of *Thysanoblatta latipennis* Kirby) British East Africa. 38. (53 BMNH). *P. dentata* Sauss. and Zehn. Pretoria, South Africa (det. Princis). 39. (989 L). *P. cribrosa* (Sauss.). Pretoria, South Africa (det. Princis). 40. (990 L). *P. thoracica* (Walker). Mamathes, Basutoland (det. Princis). 41. (991 L). *P. reflexa* (Walker). Zululand, Mtubatuba Dist., Dukuduku, South Africa (det. Princis). 42. (1406 L). *H. clavigera* (Kirby). Blouberg, Transvaal (det. Princis). (scale = Figs. 37, 41, 42 = 5 mm; Figs. 38-40 = 10 mm).



43




44



45




46




47



48




49




50


Figures 43-50. Adult males of *Hostilia*, *Blepharodera*, and *Bantua* spp.
43. (170 ANSP). *H. proterva* (Stål). Pinetown, Natal, South Africa (det. Rehn). 44. (1405 L). *H. proterva*. Stanford Hill, Durban, Natal, South Africa (det. Princis). 45. (1426 L). *Blepharodera discoidalis* (Brunner). Cap. 1872 (det. Princis). 46. (1425 L). *B. ciliata* Burm. Hopefield, C.P., X. 1883 (det. Princis). 47. (34 CUZM). *B. ciliata*. Cap. bono Spei, June 1917 (det. Princis). 48. (1404 L). *Bantua scabra* (Walker). Okahandja, S.W. Africa (det. Princis). 49. (999 L). *B. repentina* Rehn. Kruger National Park, South African Transvaal (det. Princis). 50. (1403 L). *B. robusta*. (Shelf.) Sandspruit, 15 miles northeast of Pietersburg, South African Transvaal (det. Princis). (scale = 5 mm).




Figures 51-58. Male genital phallomeres of *Hostilia* spp. 51-53. (1405 L). *H. proterva* (from ♂ shown in Fig. 44). 54-56. (170 ANSP). *H. proterva* (from ♂ shown in Fig. 43). 57-58. (1406 L). *H. clavigera* (from ♂ shown in Fig. 42). (scale = 0.2 mm).




Figures 59-70. Male genital phallomeres of *Pilema* spp. 59-61. (989 L). *P. cribrosa* (from ♂ shown in Fig. 39). 62-64. (53 BMNH). *P. dentata* (from ♂ shown in Fig. 38). 65-67. (900 L). *P. thoracica* (from ♂ shown in Fig. 40). 68-70. (991 L). *P. reflexa* (from ♂ shown in Fig. 41). (scale = 0.2 mm).



Figures 71-79. Male genital phallomeres of *Blepharodera* spp. 71-73. (1426 L). *B. discoidalis* (from ♂ shown in Fig. 45). 74-76. (34 CUZM). *B. discoidalis* (from ♂ shown in Fig. 47). 77-79. (1425 L). *B. ciliata* (from ♂ shown in Fig. 46). (scale = 0.2 mm).



Figures 80-88. Male genital phallomeres of *Bantua* spp. 80-82. (1403 L). *B. robusta* (from ♂ shown in Fig. 50). 83-85. (148 ANSP). *B. robusta* (Shelford), paratype of *Bantua armigera* Rehn. 86-88. (54 BMNH). *B. dispar*. Distant Coll. (labeled *Derocalymma pallipes* Kirby). (scale = 0.2 mm).



Figures 89-97. Male genital phallomeres of *Bantua* and *Cyrtotria* spp.
89-91. (999 L). *B. repentina* (from ♂ shown in Fig. 49). 92-94. (1404 L).
B. scabra (from ♂ shown in Fig. 48). 95-97. (993 L). *C. graniger* (from
♂ shown in Fig. 99). (scale = 0.2 mm).

In *Perisphaeria* 2 species groups can be distinguished based on the R₂:

Species Group 1. R₂ with a distinct, deep, subapical incision (Fig. 23). *P. stylifera* (Figs. 7-8, 28-33), *P. aspera* (Figs. 9, 34-36); these 2 species also have very similar shaped L_{2d}'s (Figs. 28, 34); *P. ruficornis* (Figs. 6, 22-24).

Species Group 2. R₂ without a subapical incision, or, if present, weakly represented. *P. virescens* (Figs. 2, 10-12) has a slight incision in the middle of the hook (Fig. 11) and its L_{2d} lacks any extension (Fig. 10). *P. scabra* (Figs. 3, 13-18) and *P. saxicola* (Figs. 9, 19-21) lack incisions in their R₂'s (Figs. 14, 17, 20); both species have smooth outgrowths arising from the right side of the spicular L_{2d} (Figs. 13, 16, 19).

In *Hostilia* (Figs. 42-44, 51-58) the lower margin and left side of L_{2d} is spicular and a smooth hook-like extension arises from the upper right corner (Figs. 51, 54, 57). R₂ has a shallow subapical incision (Figs. 52, 55).

In *Pilema* (Figs. 39-41, 59-70), the outer margin of R₂ is relatively straight. Three species, *cibrosa*, *dentata*, and *thoracica* have deep subapical incisions (Figs. 60, 63, 66), whereas *reflexa* (Fig. 69) has a relatively shallow one. *Thoracica* and *reflexa* have smooth outgrowths from the right side of L_{2d} (Figs. 65, 68), which appear to be lacking or weakly indicated in *cibrosa* and *dentata* (Figs. 59, 62).







In *Blepharodera* (Figs. 45-47, 71-79) the spicules on L_{2d} are very minute; the L_{2d} outgrowth is broad in *P. discoidalis* (Fig. 71) and much narrower in *B. ciliata* (Figs. 74, 77). R₂ of *discoidalis* has a minute lobe at about the middle of the hook (Fig. 72), whereas in *ciliata* there is no subapical incision (Figs. 75, 78).

Tribal Group II. Part of the L_{2d} surface on the left is reticulate (not spicular) (e.g. Figs. 80, 121), but otherwise very similar to Group I having a lateral extension on the right: *Bantua* (Figs. 48-50, 80-94) and *Cyrtotria* (Figs. 19-21, 94-144).


All of the 5 species of *Bantua* examined have R₂'s with deep subapical incisions that extend almost to the middle of the curved part of the hook (Figs. 81, 84, 87, 90, 93). Their L_{2d}'s are also basically similar in having a smooth extension on the right side arising from the reticulate portion of the sclerite (Figs. 80, 83, 86, 89, 92).

Cyrtotria is very close to *Bantua* as evidenced by the similarity of the L_{2d}'s of some of the species. There are 2 groups of *Cyrtotria* based on the presence or absence of a subapical incision:


Species Group 1. Subapical incision on R₂ absent. *C. pallicornis* (Figs. 108, 109-111), *C. marshalli* (Figs. 106, 107, 121-126), *C. latipennis* (Figs. 1, 3, 7, 101, 136-144).




Figures 98-108. Adult males of *Cyrtotria* spp. 98. (63 BMNH). *C. gibbicollis* (Stal). Natal. (labeled *Perisphaeria linearis* Walker, selected as a lectotype by K. Princis). 99. (993 L). *C. graniger* (Sauss. and Zehn.). Mudugh Prov., Somalia (det. Princis). 100. (997 L). *C. tuberculata* Shelford. Lukuledi, German East Africa (det. Princis). 101. (151 ANSP). *C. latipennis* (Kirby). N.E. Africa (det. Princis; Rehn det. the specimen as *C. capucina*, but the genitalia are similar to the type of *latipennis* (cf. Figs. 142-144 and 136-138); the L2d of *capucina* (Fig. 130) differs from that of *latipennis* (Fig. 136)). 102. (998 L). *C. orientalis* (Chopard). Kasenyi, East Congo (det. Princis). 103. (56 CUZM). *C. orientalis* Kasenyi, East Congo. 104. (996 L). *C. capucina* (Gerst.). Kakamego (det. Princis). 105. (15 HO). *C.* sp. Damaraland, South Africa. Shelford selected this specimen as the ♂ holotype of *C. poduriformis* (Walker) but Princis (personal communication) claimed that he had no authority to select an additional type to the ♀ holotype in the British Museum; the L2d surface (Fig. 25) and outgrowth differ somewhat from the L2d of the specimen which Princis believes to be *poduriformis* (Fig. 133). 106. (995 L). *C. marshalli* Shelford. Espongabera (det. Princis). 107. (14 HO). *C. marshalli*, holotype, Salisbury, Mashonaland, 5000'. 108. (992 L). *C. pallicornis* (Kirby). Pretoria, South African Transvaal (det. Princis). (scale = 5 mm).




Figures 109-117. Male genital phallomeres of *Cyrtotria* spp. 109-111. (992 L). *C. pallicornis* (from ♂ shown in Fig. 108). 112-114. (55 CUZM). *C. tuberculata*. Fsiolo, Kenya (det. Princis). 115-117. (56 CUZM). *C. orientalis* (from ♂ shown in Fig. 103). (scale = 0.1 mm).




Figures 118-126. Male genital phallomeres of *Cyrtotria* spp. 118-120. (63 BMNH). *C. gibbicollis* (from ♂ shown in Fig. 98). 121-123. (14 HO). *C. marshalli* (from ♂ shown in Fig. 107). 124-126. (995 L). *C. marshalli* from ♂ shown in Fig. 106. (scale = 0.2 mm).



Figures 127-135. Male genital phallomeres of *Cyrtotria* spp. 127-129. (998 L). *C. orientalis* (from ♂ shown in Fig. 102). 130-132. (996 L). *C. capucina* (from ♂ shown in Fig. 104). 133-135. (55 BMNH). *C. poduiformis*. Natal, South Africa (det. Princis). (scale = 0.1 mm).




Figures 136-144. Male genital phallomeres of *Cyrtotria latipennis*. 136-138. (62 BMNH). (from ♂ shown in Fig. 37). 139-141. (995 L). Kitui (det. Princis). 142-144. (151 ANSP). (from ♂ shown in Fig. 101). (scale = 0.1 mm).




145



146




147



148

Figures 145-148. Adult males of *Zuluia* spp. 145. (1408 L). *Z. lithostrota* Rehn. Letslele Valley, Gravelotte Dist., South African Transvaal (det. Princis). 146. (154 ANSP). *Z. pubescens* (Sauss.). Durban, paratype of *Z. mitella* Rehn. 147. (145 ANSP). *Z. abscissa* (Walker). Durban (det. Rehn; Princis [1960, p. 141] states that Rehn's identification of *abscissa* was an error and that the specimen was *Z. lithostrota* Rehn. The L2d of this specimen (Fig. 158) is more like that of *abscissa* (Fig. 155) than *lithostrota* (Fig. 149)). 148. (1407 L). *Z. abscissa* (det. Princis). (scale = 5 mm).



Figures 149-160. Male genital phallomeres of *Zuluia* spp. 149-151. (1408 L). *Z. lithostrota* (from ♂ shown in Fig. 145). 152-154. (154 ANSP). *Z. pubescens* (from ♂ shown in Fig. 146). 155-157. (1407 L). *Z. abscissa* (from ♂ shown in Fig. 148). 158-160. (145 ANSP). *Z. abscissa* (from ♂ shown in Fig. 147). (scale = 0.1 mm).

Species Group 2. Subapical incision on R₂ present. *C. graniger* (Figs. 95-97, 99), *C. orientalis* (Figs. 102, 103, 127-129), *C. capucina* (Figs. 104, 130-132), *C. poduriformis* (Figs. 133-135), *C. tuberculata* (Figs. 100, 112-114), *C. gibbicollis* (Figs. 98, 118-120), *C. sp.* (Figs. 25-27, 105).

Zuluiini.—L_{2d} is an unevenly outlined plate, the left side of which is scale-like or smooth; R₂ with a subapical incision. L₁ similar to *Bantua*. I place *Zuluia* (Figs. 145-160), an African genus, in this tribe. *Z. abscissa* (Figs. 155, 158) has a lobe-like extension on the left side of L_{2d} which is absent in *Z. lithostrota* (Fig. 149) and *Z. pubescens* (Fig. 152).

Rehn (1932) stated that *Zuluia* is a relative of *Derocalymma* and *Hostilia*, but in general, the relationship appeared closer to *Hostilia*. The ♂ genital phallomeres do not support this suggestion.


Derocalymmini.—L₁, which is very similar in all the species examined, has a deep cleft which may curve upwards slightly; normally the margins of the 2 lobes are close together (e.g., Figs. 187, 199), but in some preparations the phallomere was distorted by pressure of the coverslip so that the sclerotized margins were widely separated (e.g., Figs. 178, 196). The upper lobe is usually somewhat narrower and more elongate than the lower lobe, and the latter lacks setae. L_{2d} is a broad flat sclerotized plate of variable shape, slightly separated from L_{2vm}, and whose surface appears weakly wrinkled or scale-like. The R₂, generally similar in all species examined, is unmodified, lacking an apical notch or subapical incision. I place *Derocalymma* (Figs. 161-227), an African genus, in this tribe.

Based on the shape of L_{2d}, I separate 2 species groups:


Species Group 1. The L_{2d} variable in shape, but lacking a finger-like projection on the right side: *D. lampyrina* (Figs. 161, 173-178), *D. kalahari* (Figs. 163, 182-184), *D. granulata* (Figs. 164, 185-190), *D. porcellio* (Figs. 165, 191-196), *D. silphoides* (Figs. 166, 197-199), *D. rajiformis* (Figs. 167, 200-205), *D. cruralis* (Figs. 168, 206-208), *D. versicolor* (Figs. 169, 209-214), *D. sp.* (Figs. 162, 179-181).

Species Group 2. Distinct finger-like projection extending from the right side of the L_{2d}: *D. costata* (Figs. 170, 215-217), *D. pluteus* (Figs. 171, 218-223), *D. scruposa scruposa* (Figs. 172, 224-226).


Rehn (1932) in a study of *Derocalymma*, using female characters, arranged the species in what seemed to him to be a natural phylogenetic order. He considered *D. pluteus* as the least specialized of the genus (the ♀ shows no traces of caudo-lateral diverging "gut-




Figures 161-166. Adult males of *Derocalymma* spp. 161. (37 CUZM). *D. lampyrina* Gerstaecker. (det. Princis). 162. (141 ANSP). *D.* sp. German East Africa (det. by Rehn as *lampyrina* and by Princis as *versicolor*; the L2d (Fig. 141) differs from both these species (see Figs. 173, 176 and 209, 212)). 163. (1414 L). *D. kalahari* Rehn, Khomashochland, S.W. Africa (det. Princis). 164. (1413 L). *D. granulata* Sauss. and Zehn. S.W. Africa (det. Princis). 165. (1416 L). *D. porcellio* Gerstaecker. Morogoro, German East Africa (det. Princis). 166. (1411 L). *D. silphoides* Bolivar. Abachaus, Darmaland (det. Princis). (scale = 5 mm).




Figures 167-172. Adult males of *Derocalymma* spp. 167. (38 CUZM). *D. rajiformis* Rehn, Kenya (det. Princis). 168. (1410 L). *D. cruralis* (Stål). Bipindi, South Cameroon (det. Princis). 169. (1412 L). *D. versicolor* Burmeister. Southern Rhodesia (det. Princis). 170. (1417 L). *D. costata* Rehn. Cameroon (det. Princis). 171. (1409 L). *D. pluteus* Rehn. Haut-Luapula, Kansenia (det. Princis). 172. (1418 L). *D. scruposa scruposa* Rehn. Togo, Misahöhi-Mitte (det. Princis). (scale = 5 mm).




Figures 173-181. Male genital phallomeres of *Derocalymma* spp. 173-178. (1415 L). *D. lamprina*. Okavango, Runtu, S.W. Africa (det. Princis). 176-178. (37 CUZM). (from ♂ shown in Fig. 161). 179-181. (141 ANSP). *D. sp.* German East Africa (from ♂ shown in Fig. 162). (scale = 0.1 mm).




Figures 182-190. Male genital phallomeres of *Derocalymma* spp. 182-184. (1414 L). *D. kalahari*. (from ♂ shown in Fig. 163). 185-190. *D. granulata*. 185-187. (1413 L). (from ♂ shown in Fig. 164). 188-190. (143 ANSP). Damara (det. Rehn). (scale = 0.1 mm).




Figures 191-199. Male genital phallomeres of *Derocalymma* spp. 191-196. *D. porcellio*. 191-193. (1416 L). (from ♂ shown in Fig. 165). 194-196. (167 ANSP). Tendaguru, Lindi, German East Africa (det. Hebard). 197-199. (1411 L). *D. silphoides* (from ♂ shown in Fig. 166). (scale = 0.1 mm).



Figures 200-208. Male genital phallomeres of *Derocalymma* spp. 200-205. *D. rajiformis*. 200-202. (38 CUZM). (from ♂ shown in Fig. 167). 203-205. (169 ANSP). Bukama (Musée du Congo) (det. Rehn). 206-208. (1410 L). *D. cruralis* (from ♂ shown in fig. 168). (scale = 0.1 mm).



Figures 209-217. Male genital phallomeres of *Derocalymma* spp. 209-214. *D. versicolor*. 209-211. (142 ANSP). Gaberones, Bechuanaland (det. Rehn). 212-214. (1412 L). (from ♂ shown in Fig. 169). 215-217. (1417 L). *D. costata* (from ♂ shown in Fig. 170). (scale = 0.1 mm).



Figures 218-226. Male genital phallomeres of *Derocalymma* spp. 218-223. *D. pluteus*. 218-220. (157 ANSP). Paratype, Mayumbe, Congo. 221-223. (1409 L). (from ♂ shown in Fig. 171). 224-226. (1418 L). *D. scruposa scruposa* (from ♂ shown in Fig. 172). (scale = 0.1 mm).


ters", the mediocephalic portion of the disk virtually without a median carinula, and the ventro-lateral faces of the pronotum are not inflated or bullate, their surface deplanate). The L_{2d} of *pluteus* with its finger-like projection falls into group 2 along with *costata* and *scruposa*, two species which are almost at the opposite extremes of Rehn's listing. The finger-like extension could represent a "primitive" character which was lost in the species placed in group 1. However, it is also possible that the L_{2d} of *pluteus* represents a more highly evolved form with the extension developing from species which lacked it.

Perisphaerini.—Unfortunately, the spelling of this tribe differs only slightly from the *Perisphaeriini*, but Rehn (1951) used *Perisphaerus* to erect the tribe based on wing venation.


L₁ of this tribe is distinctive in that the upper lobe is considerably larger than the lower lobe, with a snoutlike projection over the cleft opening and a patch of numerous, moderately spaced setae on the lower lobe (e.g., Fig. 44). The preputial membrane is almost entirely modified to form the L_{2d} which is weakly separated from the sclerotized L_{2vm}; a small part of the preputial membrane on the right side may not be sclerotized, and minute microtrichia can be seen in this region. The hook (R₂) with (Figs. 240, 281, arrows) or without an apical notch; no other genera, here examined, have an apical notch: *Perisphaerus* (Figs. 227-259), *Pseudogloemeris* (Figs. 260-262), and *Trichoblatta* (Figs. 263-289). Three of 16 known species of *Trichoblatta* are African, the others, as well as all the species of *Perisphaerus* and *Pseudogloemeris* are Asian and adjoining islands.

The genera and species can be grouped according to the degree of development of the apical notch on R₂:


- a. Apical notch absent: *Pseudogloemeris glomeris* (Figs. 260-262), *Trichoblatta sericea* (Figs. 264, 288-289).
- b. Apical notch poorly defined: *Perisphaerus flavipes* (Figs. 257-259) (the shape of L_{2d} of this species (Fig. 257) differs from all the others of the tribe), *P. armadillo* (Figs. 233-235, 248-251), and *P. glomeriformis* (Figs. 231-232, 254-256).
- c. Apical notch well developed: *Perisphaerus contiguus* (Figs. 227, 236-238), *P. flavigornis* (Figs. 228, 239-241), *P. semilunatus* (Figs. 230, 242-244), *P. aterrimus* (Figs. 229, 245-247), *Trichoblatta magnifica* (Figs. 268, 271-273), *T. oniscina* (Figs. 267, 274-276), *T. tarsalis* (Figs. 264, 277-279), *T. nigra* (Figs. 269, 280-282), *T. guerini* (Figs. 266, 283-285).




Figures 227-235. Adult males of *Perisphaerus* spp. 227. (176 ANSP). *P. contiguus* (Sauss.). Mt. Lamington, N.E. Papua (det. Hebard). 228. (39 CUZM). *P. flavigornis* Burm. Java. 229. (983 L). *P. aterrimus* (Herbst). Kangean Island, Oengkokop (det. Princis). 230. *P. semilunatus* (Hanitsch). Sakaerat, Khorat Province, Thailand (det. Princis). 231-232. (104 MCZ). *P. glomeriformis* Lucas. Samar, Catarman, Philippine Islands (det. Princis). 232. Tooth (arrow) on underside of pronotum. 233-235. *P. armadillo* Serville. 233. (989 L). Prai Jawang, East Sumba (det. Princis). 234-235. (109 MCZ). Zamboanga, Philippine Islands (det. Princis). 235. Tooth (arrow) on underside of pronotum. (scale: Figs. 232, 235 = 1 mm; all others = 5 mm).




Figures 236-244. Male genital phallomeres of *Perisphaerus* spp. 236-238. (176 ANSP). *P. contiguus* (from ♂ shown in Fig. 227). 239-241. (39 CUZM). *P. flavicornis* (from ♂ shown in Fig. 228). 242-244. *P. semilunatus* (from ♂ shown in Fig. 230). (scale = 0.1 mm).




Figures 245-253. Male genital phallomeres of *Perisphaerus* spp. 245-247. (983 L). *P. aterrimus* (from ♂ shown in Fig. 229). 248-251. *P. armadillo*. 248-249. (109 MCZ). (from ♂ shown in Fig. 234). 250-251. (982 L). (from ♂ shown in Fig. 233). 252-253. (180 ANSP). *P. aeaneus* (Brunner). Sibuyan Island (det. Hebard). (scale = 0.1 mm).




Figures 254-262. Male genital phallomeres of *Perisphaerus* and *Pseudoglomeris* spp. 254-256. (104 MCZ). *Perisphaerus glomeriformis* (from ♂ shown in Fig. 231). 257-259. (51 BMNH). *Perisphaerus flavipes* (Sauss.). Kinta Valley, Perak (det. Princis; this specimen was det. as *Perisphaeria lucasiana* Sauss. and Zehn. (a synonym) by Hanitsch). 260-262. (46 BMNH). *Pseudoglomeris glomeris* (Sauss.). Matale, Ceylon (det. Princis).



Figures 263-270. Adult males of *Trichoblatta* spp. 263. (987 L). *T. pygmaea* (Karny). Ryukyu (det. Princis). 264. (985 L). *T. tarsalis* (Walker). Kambaiti, N.E. Burma, 7000' (det. Princis). 265. (984 L). *T. sericea* (Sauss.). Madras, India (det. Princis). 266. (8 BMNH). *T. guerini* Sauss. and Zehn. "Givalior" (det. Princis). 267. (988 L). *T. oniscina* (Gerstaecker). Urwald, Ukaika (det. Princis). 268. (986 L). *T. magnifica* (Shelford). Tonkin, Chapa, Indo-China (det. Princis). 269-270. (103 MCZ). 269. *T. nigra* (Shiraki). Formosa (det. Princis). 270. Tooth (arrow) on underside of pronotum. (scale: Figs. 263-269 = 5 mm; Fig. 270 = 1 mm).



Figures 271-279. Male genital phallomeres of *Trichoblatta* spp. 271-273. (986 L). *T. magnifica* (from ♂ shown in Fig. 268). 274-276. (988 L). *T. oniscina* (from ♂ shown in Fig. 267). 277-279. (985 L). *T. tarsalis* (from ♂ shown in Fig. 264). (scale = 0.1 mm).



Figures 280-289. Male genital phallomeres of *Trichobolatta* spp. 280-282. (103 MCZ). *T. nigra* (from ♂ shown in Fig. 269). 283-285. (8 BMNH). *T. guerini* (from ♂ shown in Fig. 266). 286-287. (987 L). *T. pygmaea* (from ♂ shown in Fig. 263). 288-289. (984 L). *T. sericea* (from ♂ shown in Fig. 265). (scale = 0.1 mm).

Some genera of *Perisphaeriinae* have a characteristic tooth (Figs. 232, 235, 270) on each side underneath the pronotum into which the apex of the tegmina fit. Princis (1960) and Beier (1961) used this structure as a characteristic of *Derocalymminae* and *Derocalymmini*, respectively. The shape of the tooth may have value as a species character (cf. Figs. 232 and 235).

Shelford (1908) summarized Saussure and Zehntner's (1895) analysis of the evolution of the form of the pronotum in the *Perisphaeriinae*. The complex type of the pronotum of genera such as *Pilema* and *Cyrtotria* appear to have evolved from a more simple form. *Pilema thoracica* has been found in family groups of a female and nymphs in round holes in the ground about 6 inches in depth. Shelford suggests that the shovel-like pronotum of *Pilema* is probably used for digging and evolved as a result of a change to this habitat. *Derocalymma* spp. live under heavy stones and their flattened form would enable them to crawl through narrow crevices for security from predators. Some species of *Cyrtotria* also have the pronotum adapted for digging, but others haven't; nothing is known of the habits of this genus. The *Bantua* pronotum is intermediate in structure between *Cyrtotria* and *Derocalymma*, but nothing is known of the genus other than it has been collected under a log. Shelford suggests that the *Derocalymma* form originated from a form like *Pilema* evolving through a *Bantua*-like stage.

SUMMARY

Based on male genitalia, 11 genera of *Perisphaeriinae* are placed in 4 tribes as follows:

1. *Perisphaeriini*: *Perisphaeria*, *Blepharodera*, *Pilema*, *Hostilia*, *Bantua*, and *Cyrtotria*.
2. *Zuluiini*: *Zulia*.
3. *Derocalymmini*: *Derocalymma*.
4. *Perisphaerini*: *Perisphaerus*, *Pseudoglomeris*, and *Trichoblatta*.






ACKNOWLEDGEMENTS

I thank the following for the loan of museum specimens: Dr. David Rentz, Academy of Natural Sciences of Philadelphia; Dr. David Ragge, British Museum (Natural History); Dr. Karl Princis, Zoological Institute, Lund, Sweden; Dr. M. W. R. de V. Graham, Hope Department Entomology, Oxford, England; Dr. Howard Evans, formerly at the Museum of Comparative Zoology, Harvard University; Dr. S. L. Tuxen, Zoological Museum, Copen-

hagen, Denmark. I am grateful to Mr. Samuel H. Cohen for taking the photographs.

REFERENCES

- BEIER, M.
- 1961. H. G. Bronns Klassen und Ordnungen des Tierreichs. Arthropoda, Insecta, Blattpteroidea, Blattodea. 5, Div. III, Book 6. Part 4, pp. 587-848. Leipzig.
- GURNEY, A. B. AND L. M. ROTH
- 1972. A generic review of the cockroaches of the subfamily Panchlorinae (Dictyoptera, Blattaria, Blaberidae). Ann. Entomol. Soc. Amer. 65: 521-532.
- McKITTRICK, F. A.
- 1964. Evolutionary studies of cockroaches. Cornell Univ. Agric. Exp. Sta. Memoir 389, 197 pp.
- PRINCIS, K.
- 1960. Zur systematik der Blattarien. Eos 36: 427-449.
 - 1963. Blattariae. In South African Animal Life. Stockholm, pp. 1-318.
- REHN, J. A. G.
- 1922. Contributions to our knowledge of the Dermaptera and Orthoptera of the Transvaal and Natal. Part 1. Dermaptera and Blattidae. Ann. Transvaal Mus. 9: 1-99.
 - 1932. African and Malagasy Blattidae (Orthoptera). Part II. Proc. Acad. Nat. Sci. Philadelphia 84: 405-511.
 - 1933. Dermaptera and Orthoptera of the De Schauensee South African Expedition, — Part 1. Proc. Acad. Nat. Sci. of Philadelphia 85: 39-66.
- REHN, J. W. H.
- 1951. Classification of the Blattaria as indicated by their wings (Orthoptera). Mem. Amer. Entomol. Soc. No. 14: 1-134.
- ROTH, L. M.
- 1970. The male genitalia of Blattaria. III. Blaberidae: Zetoborinae. Psyche 77: 217-236.
 - 1970a. The male genitalia of Blattaria. IV. Blaberidae: Blaberinae. Psyche 77: 308-342.
 - 1971. The male genitalia of Blattaria. VIII. *Panchlora*, *Anchoblatta*, *Bolleya*, *Pelloblatta*, and *Achroblatta* (Blaberidae: Panchlorinae). Psyche 78: 296-305.
 - 1972. The male genitalia of Blattaria. IX. Blaberidae. *Gyna* spp. (Perisphaeriinae), *Phoraspis*, *Thorax*, and *Phlebonotus* (Epilamprinae). Psyche 98: 185-217.
 - 1973. *Paramuzoa* (Nyctiborinae), a new cockroach genus previously confused with *Parasphaeria* (Epilamprinae). Psyche 80: 249-264.
- SAUSSURE, H. DE AND L. ZEHNTNER.
- 1895. Revision de la Tribu des Perisphaeriens. Insectes Orthoptères de la famille des Blattides. Rev. suisse Zool. 3: 1-59.
- SHELFORD, R.
- 1908. Some new genera and species of Blattidae, with notes on the form of the pronotum in the subfamily Perisphaeriinae. Ann. and Mag. Hist. Ser. 8, 1: 157-177.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

